


Infant Mortality

Indiana 2022



Division of **Maternal & Child Health**



Infant Mortality Rates



Infant Mortality Facts

- Infant mortality is defined as the death of a baby before his or her first birthday
- The infant mortality rate (IMR) is the number of babies who die in the first year of life per 1,000 live births
- In 2022, there were 577 infant deaths across the state, making the infant mortality rate 7.2
- In Indiana, the Non-Hispanic (NH) Black infant mortality rate was more than double the NH White rate
- Neonatal Infant Death = 0 – 27 days
- Post-Neonatal Infant Death = 28 – 364 days

Infant Mortality Rates by Age of Death, 2022




2022	
HOSPITAL REGION	IMR
CENTRAL	8.1
CENTRAL SOUTHWESTERN	4.6*
EASTERN	7.2
MIDWESTERN	5.5*
NORTHEASTERN	7.5
NORTHERN	7.5
NORTHWESTERN	6.6
SOUTHEASTERN	5.2*
SOUTHERN	5.7
SOUTHWESTERN	6.9
WESTERN	8.3

* Rates based on counts less than 20 are considered unstable and should be interpreted with caution.

How to Reduce Infant Mortality

- Improve overall health for women of child-bearing age
- Promote early and adequate prenatal care
- Decrease early elective deliveries before 39 weeks
- Decrease prenatal smoking and substance use
- Increase breastfeeding duration and exclusivity
- Support birth spacing and interconception wellness
- Promote the ABCs of safe sleep: place baby to sleep alone, on his or her back, in a crib

Causes of Infant Mortality Indiana 2022



Birth Outcome Indicators

Indiana 2022



Division of
Maternal & Child Health


Indiana Quick Facts

- 79,675 live births:
 - 68.9% NH White
 - 12.5% NH Black
 - 12.4% Hispanic
- Similar low birthweight (LBW) and preterm rate as the nation
- Similar percentage of mothers on Medicaid as the nation
- Nearly 1/3 of pregnant women do not receive early prenatal care (PNC)

Disparities in Indiana

- NH Black infants in Indiana are more likely to be born LBW or preterm
- NH Black and Hispanic women are more likely to not receive early prenatal care (1st trimester)
- NH Black women in Indiana are less likely to breastfeed at hospital discharge
- NH White women in Indiana are more likely to smoke during pregnancy

Birth Outcome Indicators by Race & Ethnicity, 2022



LBW = Low Birthweight PNC = Prenatal Care IMR = Infant Mortality Rate NH = Non-Hispanic

2022 REGION	% LBW (< 2,500 G)	% PRETERM (<37 WKS GESTATION)	% NO EARLY PNC (1 ST TRIMESTER)	% ~SMOKING	% NOT BREASTFEEDING	% MOTHER ON MEDICAID
NORTHWESTERN	8.6	11.2	32.7	3.6	18.6	46.6
NORTHERN	8.2	10.6	36.2	6.0	15.1	46.8
NORTHEASTERN	8.1	9.5	34.2	6.2	13.7	33.3
MIDWESTERN	8.3	10.5	27.9	11.2	20.1	48.2
WESTERN	8.9	10.4	23.7	6.6	13.4	33.5
EASTERN	9.9	11.8	23.5	11.8	24.8	49.1
CENTRAL SOUTHWESTERN	8.5	10.6	25.3	11.1	19.3	45.0
CENTRAL	9.2	11.3	29.4	4.2	12.3	39.2
SOUTHEASTERN	7.8	11.0	29.3	11.1	19.4	42.0
SOUTHWESTERN	8.7	11.0	21.2	7.4	22.1	42.2
SOUTHERN	8.5	11.4	24.2	10.3	19.8	42.2
INDIANA	8.7	10.9	29.1	6.6	16.1	41.1
U.S.	8.6	10.4	23.0	3.7	14.8	41.3

~Due to a change in reporting methods, smoking during pregnancy should not be compared to previous years data.

For more information on infant mortality, visit:
www.in.gov/health/mch/data/infant-mortality

Created by Maternal and Child Health Epidemiology, February 2024

