


INFANT MORTALITY

Southeastern Hospital Region 2020

For more information on infant mortality in your area, please see the Indiana Department of Health Stats Explorer, Mortality Data (https://gis.in.gov/apps/isdh/meta/stats_layers.htm)




Southeastern District
IMR = 6.1*
N = 17



INFANT MORTALITY FACTS


- Infant mortality is defined as the death of a baby before his or her first birthday.
- The infant mortality rate (IMR) is the number of babies who die in the first year of life, per 1,000 live births.
- Of the 522 Indiana infant deaths in 2020, 17 occurred in the Southeastern Hospital Region.
- Non-Hispanic black infants are 2.4 times more likely to die than Non-Hispanic white infants in Indiana.

Infant Mortality Rates by Age at Death




Note: Age-specific and cause-specific mortality rates may not sum to the overall infant mortality rate due to rounding.

Infant Mortality Rates by Cause of Death



* Numerator less than 20, rate unstable
** Rate suppressed due to low counts

Infant Mortality Rates 2016 - 2020



	2016	2017	2018	2019	2020
—●— Southeastern Region	7.2	7.8	5.1*	7.1	6.1*
—■— Indiana	7.5	7.3	6.8	6.5	6.6
—▲— U.S.	5.9	5.8	5.7	5.6	5.4
—■— HP 2020 Goal	6.0	6.0	6.0	6.0	6.0

PRACTICES TO REDUCE INFANT MORTALITY

- Improve overall health for women of child-bearing age.
- Promote early & adequate prenatal care.
- Decrease early elective deliveries before 39 weeks.
- Decrease prenatal smoking & substance use.
- Increase breastfeeding duration & exclusivity.
- Support birth spacing & interconception wellness.
- Promote the ABCs of safe sleep: place baby to sleep alone, on his or her back, in a crib.

BIRTH OUTCOME INDICATORS

Southeastern Hospital Region 2020

For more information on birth outcomes in your area, please see the Indiana Department of Health Stats Explorer, Natality Data (https://gis.in.gov/apps/isdh/meta/stats_layers.htm)


INDIANA QUICK FACTS

- Smoking rates among pregnant women continue to be much higher than the national rate.
- Almost 1/3 of pregnant women do not receive early PNC.
- NH black babies in Indiana are more likely to be LBW or preterm.
- Hispanic and NH black women in Indiana are more likely to not receive early 1st trimester PNC.
- NH white women in Indiana are more likely to smoke during pregnancy.

SOUTHEASTERN REGION QUICK FACTS

- There were 2,792 births in the Southeastern Region in 2020.
- Lower percentages of LBW and preterm births compared to the state of Indiana.
- Higher percentage of women receiving early PNC.
- Higher percentage of women smoking during pregnancy.
- Lower percentage of breastfeeding at hospital discharge.

Outcomes by Race and Ethnicity, Southeastern Region, 2020



LBW = Low Birthweight PNC = Prenatal Care IMR = Infant Mortality Rate NH = Non-Hispanic

	% LBW ($< 2,500$ G)	% PRETERM (< 37 WKS GESTATION)	% NO EARLY PNC (1 ST TRIMESTER)	% SMOKING	% NOT BREASTFEEDING	% MOTHER ON MEDICAID
DEARBORN	5.7	8.6	24.0	14.1	29.5 ^S	33.9
DECATUR	4.9*	8.5	22.3	17.0	25.9 ^S	41.6
FAYETTE	7.1*	6.7*	30.5	20.9 ^S	31.4 ^S	52.7 ^S
FRANKLIN	8.0*	11.0	30.8	13.9	23.6	27.0 ^S
OHIO	**	11.3*	24.5*	15.1*	43.4 ^S	35.8*
RIPLEY	5.7	9.3	29.2	14.5	28.4 ^S	34.7
RUSH	6.3*	9.5*	20.0	22.1 ^S	28.9 ^S	46.8
UNION	**	**	21.6*	14.9*	23.0*	25.7*
WAYNE	9.8	12.7	30.6	17.6 ^S	24.5	53.8
SOUTHEASTERN REGION	7.1	9.9	27.2	16.6	27.3	42.3
INDIANA	8.1	10.4	30.7	10.9	18.0	38.5

* = Fewer than 20 birth outcomes, percentage unstable

** = Fewer than 5 birth outcomes, rate suppressed

S = County significantly different from state