

Table 3A
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution ⁽²⁾								
Clinic	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaska Native	Asian/Pacific Islander	Multiple Races	All Other/Unknown
		%	%	%	%	%	%	%
Action Ctr	40	*	*	*	*	*	*	*
Albion	304	79.6	0.0	18.8	0.0	1.0	0.7	0.0
Anderson	866	76.8	13.3	7.9	0.2	0.1	1.2	0.6
Bartholome	526	81.9	1.7	14.3	0.0	0.6	0.8	0.8
Bedford	285	96.1	0.0	1.1	0.0	1.1	0.4	1.4
Benton Co	45	*	*	*	*	*	*	*
Berne	38	*	*	*	*	*	*	*
Bicknell	30	*	*	*	*	*	*	*
Blackburn	454	18.7	70.7	6.2	0.2	1.5	2.6	0.0
Blackford	102	95.1	1.0	3.9	0.0	0.0	0.0	0.0
Boone Co	202	92.6	3.0	2.5	0.0	1.0	0.0	1.0
Brazil	231	95.2	1.3	0.4	0.0	0.4	1.7	0.9
Brown Co	64	*	*	*	*	*	*	*
Cambridge	28	*	*	*	*	*	*	*
Carroll Co	82	*	*	*	*	*	*	*
Cass Co	384	64.6	1.0	31.0	0.0	0.8	1.3	1.3
Citizens	654	40.4	46.0	10.7	0.0	1.1	1.4	0.5
Civic Ctr	1,021	73.5	16.7	5.8	0.3	0.8	2.3	0.7
Clark Co	722	71.6	9.8	13.6	0.4	1.1	2.8	0.7
Clay City	5	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3A
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution ⁽²⁾								
Clinic	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaska Native	Asian/Pacific Islander	Multiple Races	All Other/Unknown
		%	%	%	%	%	%	%
Clinton	89	*	*	*	*	*	*	*
Clinton Co	269	62.5	0.4	34.9	0.4	0.0	0.7	1.1
Comm East	119	33.6	59.7	4.2	0.0	0.0	2.5	0.0
Cottage Gr	864	70.7	12.4	14.7	0.0	0.3	1.4	0.5
Crawford	82	*	*	*	*	*	*	*
Crown Pt	558	81.7	6.3	9.0	0.0	1.8	0.2	1.1
Daviess Co	196	77.6	1.0	17.9	0.0	0.0	2.0	1.5
Dearborn	330	96.7	0.9	0.3	0.3	0.0	0.6	1.2
Decatur	161	98.1	0.0	1.2	0.0	0.0	0.0	0.6
Decatur Co	194	97.9	0.0	1.5	0.5	0.0	0.0	0.0
Dekalb Co	306	93.8	0.0	4.2	0.0	0.0	1.3	0.7
Dubois Co	252	75.4	0.0	21.0	0.0	2.0	1.2	0.4
E Chicago	568	6.9	37.9	53.9	0.2	0.4	0.4	0.5
Eaton	11	*	*	*	*	*	*	*
Elkhart	1,321	55.4	13.7	28.5	0.0	0.8	1.0	0.6
Elwood	91	*	*	*	*	*	*	*
Fayette Co	189	96.3	2.1	0.5	0.0	0.5	0.0	0.5
Floyd Co	476	77.3	13.0	5.0	0.2	1.9	1.7	0.8
Forest Man	719	10.4	81.8	4.2	0.3	0.8	1.5	1.0
Fountain	111	94.6	0.0	3.6	0.0	0.9	0.0	0.9

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3A
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution ⁽²⁾								
Clinic	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaska Native	Asian/Pacific Islander	Multiple Races	All Other/Unknown
		%	%	%	%	%	%	%
Franklin C	100	98.0	0.0	1.0	0.0	0.0	0.0	1.0
Fulton Co	143	90.2	0.0	9.1	0.0	0.0	0.7	0.0
Georgetown	1,402	19.4	45.9	31.2	0.3	1.1	1.1	0.9
Gibson Co	200	93.5	2.5	1.5	1.0	0.5	0.5	0.5
Glenn Pk	594	16.8	67.0	12.1	0.3	0.8	2.0	0.8
Goshen	622	56.8	1.4	39.1	0.0	1.1	1.0	0.6
Grant Co	564	77.1	13.5	5.5	0.0	0.5	2.1	1.2
Grant St	415	2.2	89.9	5.8	0.2	0.2	1.0	0.7
Grassy Crk	965	21.9	56.1	18.9	0.0	0.7	2.0	0.5
Greene Co	164	97.0	0.0	1.8	0.0	0.0	0.6	0.6
Hamilton	682	78.3	7.2	10.4	0.1	2.5	1.3	0.1
Hammond	669	26.6	31.2	38.0	0.0	0.6	1.9	1.6
Hancock Co	296	94.9	1.4	2.4	0.0	0.0	1.0	0.3
Hanna	1,393	59.1	6.5	14.3	0.2	18.0	0.9	1.0
Harrison	254	90.9	0.4	5.1	0.0	0.0	2.0	1.6
Hendricks	421	85.3	6.2	3.3	0.0	3.8	0.0	1.4
Henry Co	300	93.3	1.0	3.0	0.3	0.3	1.3	0.7
Hessville	364	53.0	15.7	29.7	0.0	0.5	0.8	0.3
Hispanic C	133	3.8	13.5	79.7	0.0	0.8	0.0	2.3
Howard Co	705	85.1	8.1	3.7	0.0	0.3	1.8	1.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3A
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution ⁽²⁾								
Clinic	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaska Native	Asian/Pacific Islander	Multiple Races	All Other/Unknown
		%	%	%	%	%	%	%
Huntington	232	96.1	0.4	1.7	0.0	0.0	1.3	0.4
Jackson Co	352	84.1	1.1	13.9	0.0	0.0	0.0	0.9
Jasonville	7	*	*	*	*	*	*	*
Jasper Co	217	85.7	0.9	12.4	0.0	0.0	0.5	0.5
Jay Co	158	90.5	0.6	6.3	0.0	0.0	0.0	2.5
Jefferson	224	85.7	1.3	7.6	0.0	0.9	2.2	2.2
Jennings	284	95.4	0.7	2.1	0.7	0.4	0.7	0.0
Johnson Co	730	87.8	2.1	4.9	0.0	2.6	1.2	1.4
Kosciusko	441	77.6	1.1	20.0	0.0	0.5	0.0	0.9
Lagrange	145	83.4	1.4	12.4	0.0	0.7	0.0	2.1
Lake Station	512	50.2	19.1	26.8	0.0	0.6	3.1	0.2
Laporte	423	81.1	2.6	13.5	0.0	0.5	1.4	0.9
Main St	837	83.2	11.5	2.0	0.1	0.6	2.2	0.5
Marshall	373	76.4	0.3	22.0	0.0	0.3	1.1	0.0
Martin Co	57	*	*	*	*	*	*	*
Memorial	183	37.2	29.5	26.8	0.0	3.8	2.2	0.5
Merrillville	2	*	*	*	*	*	*	*
Methodist	225	30.2	52.4	13.3	0.0	3.6	0.4	0.0
Miami Co	202	90.6	2.5	3.0	0.0	0.5	3.5	0.0
Mich City	455	56.0	34.5	4.8	0.0	1.1	2.9	0.7

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3A
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution ⁽²⁾								
Clinic	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaska Native	Asian/Pacific Islander	Multiple Races	All Other/Unknown
		%	%	%	%	%	%	%
Moms/Kids	125	30.4	33.6	32.8	0.0	0.8	1.6	0.8
Monroe Co	657	82.3	4.7	3.8	0.0	6.4	2.4	0.3
Montgomery	282	86.9	1.4	8.5	0.4	0.7	1.1	1.1
Morgan Co	423	96.9	0.7	1.7	0.0	0.0	0.2	0.5
N Arlingto	838	33.7	46.4	15.3	0.0	1.1	3.0	0.6
N Liberty	18	*	*	*	*	*	*	*
N Manchest	52	*	*	*	*	*	*	*
Newton Co	57	*	*	*	*	*	*	*
Northwest	624	16.2	41.8	37.7	0.3	2.4	1.1	0.5
Ohio Co	7	*	*	*	*	*	*	*
Orange Co	159	95.6	1.9	0.0	0.0	1.3	0.6	0.6
Owen Co	116	94.8	0.9	2.6	0.0	0.9	0.9	0.0
Parke Co	75	*	*	*	*	*	*	*
Perry Co	127	99.2	0.0	0.0	0.0	0.0	0.0	0.8
Pike Co	74	*	*	*	*	*	*	*
Portage	404	71.3	10.9	15.8	0.0	0.2	0.0	1.7
Posey Co	98	*	*	*	*	*	*	*
Pulaski Co	70	*	*	*	*	*	*	*
Purdue	133	30.1	6.8	6.8	0.0	54.1	0.8	1.5
Putnam Co	222	94.6	1.4	1.8	0.0	0.9	0.5	0.9

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3A
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution ⁽²⁾								
Clinic	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaska Native	Asian/Pacific Islander	Multiple Races	All Other/Unknown
		%	%	%	%	%	%	%
Randolph	190	86.8	0.0	8.4	0.5	0.5	1.6	2.1
Richmond	577	81.6	6.8	6.9	0.3	0.9	2.4	1.0
Ripley Co	196	93.9	0.0	2.0	0.5	0.0	1.5	2.0
Rush Co	106	97.2	0.0	0.9	0.0	0.9	0.0	0.9
SB Ireland Rd	711	70.0	17.6	7.6	0.3	1.5	2.4	0.6
SB Lafayette	999	34.5	37.6	20.8	0.4	1.6	2.9	2.1
Scott Co	227	97.8	0.0	1.3	0.0	0.0	0.4	0.4
Shelby Co	324	87.0	1.9	9.9	0.0	0.0	0.6	0.6
Shelters	4	*	*	*	*	*	*	*
SJ Reg Ctr	47	*	*	*	*	*	*	*
So Calhoun	2,409	46.4	28.9	15.7	0.1	6.0	2.2	0.8
Spencer Co	102	89.2	1.0	5.9	0.0	0.0	0.0	3.9
St Francis	1	*	*	*	*	*	*	*
St V-FLC	142	21.1	62.0	15.5	0.0	0.0	1.4	0.0
Starke Co	171	96.5	0.0	2.3	0.0	0.6	0.0	0.6
Steuben Co	275	89.5	0.7	6.2	0.0	1.1	0.4	2.2
Sullivan	158	98.7	0.0	1.3	0.0	0.0	0.0	0.0
Switz""Land	44	*	*	*	*	*	*	*
Tippecanoe	1,140	65.2	12.0	19.5	0.3	0.8	1.1	1.1
Tipton Co	61	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3A
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution ⁽²⁾								
Clinic	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaska Native	Asian/Pacific Islander	Multiple Races	All Other/Unknown
		%	%	%	%	%	%	%
Union Co	48	*	*	*	*	*	*	*
United Fai	342	78.9	11.1	5.6	0.0	0.3	2.9	1.2
Valpo	382	84.3	5.2	7.1	0.0	1.8	0.3	1.3
Vigo	676	83.9	10.8	2.4	0.0	0.9	1.2	0.9
Vincennes	209	94.3	0.5	1.0	0.0	0.5	1.4	2.4
W 21st Ave	354	2.5	96.0	0.8	0.0	0.0	0.3	0.3
Wabash	165	90.3	1.2	1.2	0.0	0.6	6.1	0.6
Warren Co	67	*	*	*	*	*	*	*
Warrick Co	246	95.5	2.4	0.4	0.0	0.4	1.2	0.0
Washington	168	98.8	0.0	0.6	0.0	0.0	0.0	0.6
Wells Co	158	89.2	1.3	7.6	0.0	0.0	1.3	0.6
Westside	709	43.6	21.6	31.5	0.1	1.0	1.3	1.0
White Co	213	79.8	0.0	18.3	0.0	0.5	1.4	0.0
Whitley Co	184	97.3	0.5	1.1	0.0	0.5	0.5	0.0
Wishard	124	16.9	44.4	35.5	0.0	1.6	1.6	0.0
Women""s	1	*	*	*	*	*	*	*
Unknown Clinics	307	68.7	17.3	9.1	0.0	2.0	1.0	2.0
Indiana	46,232	64.1	18.3	13.4	0.1	1.9	1.4	0.8

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3A
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution ⁽²⁾								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Nation (Prior Year)	1,162,961	41.6	22.7	29.2	1.1	3.4	1.3	0.7

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4A
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Age Distribution by Clinic

Comparison of Age Distribution ⁽²⁾								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Action Ctr	40	*	*	*	*	*	*	*
Albion	304	0.0	6.3	14.1	63.5	15.1	1.0	0.0
Anderson	866	0.2	6.2	15.1	61.4	15.5	1.5	0.0
Bartholome	526	0.0	9.5	16.3	57.6	14.8	1.7	0.0
Bedford	285	0.0	9.8	15.4	58.9	14.4	1.4	0.0
Benton Co	45	*	*	*	*	*	*	*
Berne	38	*	*	*	*	*	*	*
Bicknell	30	*	*	*	*	*	*	*
Blackburn	454	0.2	4.8	9.5	64.3	19.2	2.0	0.0
Blackford	102	0.0	6.9	16.7	66.7	8.8	1.0	0.0
Boone Co	202	0.0	5.0	15.3	67.3	11.4	1.0	0.0
Brazil	231	0.4	7.8	16.5	63.2	10.8	1.3	0.0
Brown Co	64	*	*	*	*	*	*	*
Cambridge	28	*	*	*	*	*	*	*
Carroll Co	82	*	*	*	*	*	*	*
Cass Co	384	0.8	6.3	14.1	61.2	16.7	1.0	0.0
Citizens	654	0.0	5.7	11.3	63.0	19.0	0.9	0.2
Civic Ctr	1,021	0.2	6.8	12.5	64.9	14.7	0.8	0.1
Clark Co	722	0.1	4.4	13.2	62.3	19.1	0.7	0.1

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4A
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Age Distribution by Clinic

Comparison of Age Distribution ⁽²⁾								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Clay City	5	*	*	*	*	*	*	*
Clinton	89	*	*	*	*	*	*	*
Clinton Co	269	0.0	4.5	14.1	64.7	14.9	1.9	0.0
Comm East	119	0.0	3.4	14.3	68.9	12.6	0.8	0.0
Cottage Gr	864	0.5	5.1	12.7	63.4	17.5	0.8	0.0
Crawford	82	*	*	*	*	*	*	*
Crown Pt	558	0.0	3.0	9.5	61.3	24.9	1.1	0.2
Daviess Co	196	0.0	4.6	12.8	66.8	13.3	0.0	2.6
Dearborn	330	0.6	4.8	13.0	63.0	17.3	0.9	0.3
Decatur	161	0.0	4.3	13.7	68.9	12.4	0.6	0.0
Decatur Co	194	0.0	7.2	18.6	61.3	12.4	0.5	0.0
Dekalb Co	306	0.3	6.2	18.0	62.4	12.4	0.7	0.0
Dubois Co	252	0.0	2.8	9.9	64.7	20.2	2.4	0.0
E Chicago	568	1.1	5.5	12.9	57.4	21.1	1.9	0.2
Eaton	11	*	*	*	*	*	*	*
Elkhart	1,321	0.3	4.5	12.9	61.4	19.5	1.3	0.0
Elwood	91	*	*	*	*	*	*	*
Fayette Co	189	0.5	6.9	15.3	67.2	10.1	0.0	0.0
Floyd Co	476	0.4	6.3	16.2	62.4	13.9	0.6	0.2

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4A
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Age Distribution by Clinic

Comparison of Age Distribution ⁽²⁾								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Forest Man	719	0.4	5.6	13.6	64.0	14.9	1.5	0.0
Fountain	111	0.0	4.5	17.1	59.5	18.9	0.0	0.0
Franklin C	100	0.0	9.0	12.0	62.0	16.0	1.0	0.0
Fulton Co	143	0.0	3.5	12.6	71.3	11.2	1.4	0.0
Georgetown	1,402	0.1	4.1	10.3	58.1	25.7	1.4	0.3
Gibson Co	200	0.0	3.0	17.5	65.0	13.5	0.5	0.5
Glenn Pk	594	0.5	4.0	12.0	58.8	23.6	1.2	0.0
Goshen	622	0.5	4.2	15.0	55.6	23.2	1.6	0.0
Grant Co	564	1.1	5.9	15.1	64.5	12.8	0.7	0.0
Grant St	415	0.0	7.7	14.2	58.1	19.0	0.7	0.2
Grassy Crk	965	0.7	5.1	14.3	59.9	18.0	1.9	0.1
Greene Co	164	0.6	8.5	16.5	64.6	9.8	0.0	0.0
Hamilton	682	0.3	2.2	12.3	60.4	22.3	1.5	1.0
Hammond	669	0.3	5.2	12.0	57.0	23.0	2.5	0.0
Hancock Co	296	0.0	5.1	13.5	61.5	18.9	1.0	0.0
Hanna	1,393	0.2	4.7	8.0	65.5	20.1	1.3	0.1
Harrison	254	0.0	3.5	14.6	65.4	15.4	0.8	0.4
Hendricks	421	0.0	4.3	14.5	60.1	19.5	1.7	0.0
Henry Co	300	0.3	6.0	14.0	65.3	13.3	0.7	0.3

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4A
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Age Distribution by Clinic

Comparison of Age Distribution ⁽²⁾								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Hessville	364	0.3	6.0	12.9	53.8	25.5	1.1	0.3
Hispanic C	133	0.8	3.0	4.5	63.9	27.1	0.8	0.0
Howard Co	705	0.4	5.7	15.5	62.7	14.8	0.9	0.1
Huntington	232	0.4	5.2	20.7	62.5	9.5	0.9	0.9
Jackson Co	352	0.0	8.0	15.1	63.6	12.5	0.9	0.0
Jasonville	7	*	*	*	*	*	*	*
Jasper Co	217	0.0	5.1	15.7	55.8	20.3	3.2	0.0
Jay Co	158	0.0	7.6	10.1	68.4	13.9	0.0	0.0
Jefferson	224	0.4	7.1	14.7	62.1	14.7	0.9	0.0
Jennings	284	0.4	7.0	17.6	62.7	12.0	0.4	0.0
Johnson Co	730	0.1	5.1	15.1	63.0	14.8	1.9	0.0
Kosciusko	441	0.5	3.9	15.2	59.0	20.0	1.6	0.0
Lagrange	145	0.7	6.2	16.6	56.6	17.9	0.7	1.4
Lake Station	512	0.2	5.5	10.4	57.8	25.0	1.0	0.2
Laporte	423	0.2	3.1	12.3	64.8	18.4	1.2	0.0
Main St	837	0.1	5.1	13.3	65.4	15.3	0.8	0.0
Marshall	373	0.3	3.8	12.1	59.8	22.5	1.3	0.3
Martin Co	57	*	*	*	*	*	*	*
Memorial	183	0.5	4.9	10.9	62.3	19.1	2.2	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4A
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Age Distribution by Clinic

Comparison of Age Distribution ⁽²⁾								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Merrillville	2	*	*	*	*	*	*	*
Methodist	225	0.4	4.0	13.3	67.6	14.2	0.4	0.0
Miami Co	202	0.5	5.0	16.3	61.4	16.3	0.0	0.5
Mich City	455	0.4	4.6	14.9	58.2	19.8	2.0	0.0
Moms/Kids	125	0.0	3.2	14.4	53.6	24.8	4.0	0.0
Monroe Co	657	0.0	2.1	13.1	63.6	19.3	1.2	0.6
Montgomery	282	0.0	7.8	14.2	60.6	16.0	1.4	0.0
Morgan Co	423	0.0	5.0	11.3	67.4	15.1	0.9	0.2
N Arlingto	838	0.6	4.7	9.9	64.1	19.6	1.2	0.0
N Liberty	18	*	*	*	*	*	*	*
N Manchest	52	*	*	*	*	*	*	*
Newton Co	57	*	*	*	*	*	*	*
Northwest	624	0.0	2.9	7.9	60.9	26.4	1.9	0.0
Ohio Co	7	*	*	*	*	*	*	*
Orange Co	159	0.6	3.8	14.5	60.4	20.1	0.6	0.0
Owen Co	116	0.0	2.6	19.8	64.7	12.9	0.0	0.0
Parke Co	75	*	*	*	*	*	*	*
Perry Co	127	0.8	10.2	17.3	63.0	7.9	0.8	0.0
Pike Co	74	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4A
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Age Distribution by Clinic

Comparison of Age Distribution ⁽²⁾								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Portage	404	0.0	4.5	9.9	65.6	18.3	1.5	0.2
Posey Co	98	*	*	*	*	*	*	*
Pulaski Co	70	*	*	*	*	*	*	*
Purdue	133	0.0	0.0	3.8	59.4	35.3	1.5	0.0
Putnam Co	222	0.5	5.9	13.5	62.6	16.7	0.9	0.0
Randolph	190	0.5	5.8	15.3	61.6	15.3	1.1	0.5
Richmond	577	0.0	2.9	14.9	64.8	17.0	0.3	0.0
Ripley Co	196	0.0	4.1	13.8	68.4	12.8	1.0	0.0
Rush Co	106	0.0	10.4	19.8	61.3	7.5	0.9	0.0
SB Ireland Rd	711	0.0	6.3	12.9	58.9	20.8	1.0	0.0
SB Lafayette	999	0.1	7.1	13.7	57.1	21.0	1.0	0.0
Scott Co	227	0.4	7.0	15.4	67.8	7.9	1.3	0.0
Shelby Co	324	0.3	8.3	12.7	64.8	12.7	1.2	0.0
Shelters	4	*	*	*	*	*	*	*
SJ Reg Ctr	47	*	*	*	*	*	*	*
So Calhoun	2,409	0.2	3.7	12.6	62.2	19.5	1.3	0.4
Spencer Co	102	0.0	6.9	13.7	54.9	21.6	2.9	0.0
St Francis	1	*	*	*	*	*	*	*
St V-FLC	142	0.0	4.2	10.6	67.6	16.9	0.7	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4A
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Age Distribution by Clinic

Comparison of Age Distribution ⁽²⁾								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Starke Co	171	1.2	2.9	10.5	64.9	18.1	1.8	0.6
Steuben Co	275	0.0	5.5	19.3	63.6	11.3	0.4	0.0
Sullivan	158	0.0	5.1	22.8	59.5	12.7	0.0	0.0
Switz""Land	44	*	*	*	*	*	*	*
Tippecanoe	1,140	0.1	5.0	13.9	60.6	18.8	1.6	0.1
Tipton Co	61	*	*	*	*	*	*	*
Union Co	48	*	*	*	*	*	*	*
United Fai	342	0.0	3.2	8.8	65.5	21.3	0.9	0.3
Valpo	382	0.3	5.2	13.4	58.9	20.2	1.8	0.3
Vigo	676	0.3	8.4	16.1	63.2	11.2	0.7	0.0
Vincennes	209	0.5	2.9	17.2	65.6	13.9	0.0	0.0
W 21st Ave	354	1.1	7.3	12.7	58.5	19.5	0.6	0.3
Wabash	165	0.0	4.8	14.5	66.7	10.9	1.8	1.2
Warren Co	67	*	*	*	*	*	*	*
Warrick Co	246	0.0	7.3	15.4	65.9	10.6	0.0	0.8
Washington	168	0.0	5.4	19.0	62.5	11.9	0.6	0.6
Wells Co	158	0.6	1.9	20.3	64.6	12.7	0.0	0.0
Westside	709	0.7	5.4	9.4	63.6	20.0	0.8	0.0
White Co	213	0.5	6.1	14.6	59.2	18.3	1.4	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4A
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Age Distribution by Clinic

Comparison of Age Distribution ⁽²⁾								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Whitley Co	184	0.0	3.3	11.4	68.5	15.8	1.1	0.0
Wishard	124	0.8	5.6	8.9	58.1	23.4	3.2	0.0
Women""s	1	*	*	*	*	*	*	*
Unknown Clinics	307	0.3	6.5	11.7	65.8	14.3	1.3	0.0
Indiana	46,232	0.3	5.2	13.2	62.1	17.9	1.2	0.1
Nation (Prior Year)	1,162,961	0.3	5.3	11.5	59.4	21.7	1.7	0.2

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators ^(2, 3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy	Hypertension During Pregnancy		
	Under- weight	Over- weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)			Number	% (Rank)
Action Ctr	29	* *	* *	25	* *	* *	8	* *	7	* *	36	* *	36	* *
Albion	250	3.2 (8)	57.6 (67)	235	19.6 (57)	54.9 (64)	55	* *	49	* *	291	0.0 (1)	291	1.7(28)
Anderson	664	3.9 (20)	57.4 (66)	632	18.2 (36)	53.0 (44)	135	51.9 (21)	148	37.2(26)	841	1.5 (20)	841	1.5(21)
Bartholome	424	7.3 (90)	53.8 (35)	412	18.9 (47)	54.4 (59)	69	* *	70	* *	516	4.7 (82)	516	7.4(90)
Bedford	237	6.8 (82)	54.9 (47)	233	16.3 (15)	53.2 (46)	33	* *	51	* *	283	5.3 (92)	283	8.1(95)
Benton Co	36	* *	* *	34	* *	* *	7	* *	8	* *	43	* *	43	* *
Berne	32	* *	* *	31	* *	* *	10	* *	8	* *	38	* *	38	* *
Bicknell	26	* *	* *	22	* *	* *	7	* *	17	* *	26	* *	26	* *
Blackburn	339	5.0 (42)	56.0 (58)	317	18.6 (43)	53.0 (44)	83	* *	80	* *	437	5.5 (93)	437	8.2(96)
Blackford	80	* *	* *	76	* *	* *	22	* *	20	* *	99	* *	99	* *
Boone Co	158	5.1 (47)	50.0 (11)	155	14.8 (6)	54.2 (58)	30	* *	36	* *	198	2.0 (28)	198	0.5(4)
Brazil	196	6.6 (79)	57.1 (64)	188	22.3 (81)	48.4 (11)	27	* *	49	* *	225	3.1 (49)	225	2.7(41)
Brown Co	50	* *	* *	48	* *	* *	8	* *	18	* *	62	* *	62	* *
Cambridge	20	* *	* *	20	* *	* *	3	* *	6	* *	28	* *	28	* *
Carroll Co	66	* *	* *	63	* *	* *	10	* *	16	* *	81	* *	81	* *
Cass Co	326	4.3 (27)	55.5 (51)	311	22.5 (86)	47.9 (9)	61	* *	36	* *	372	3.0 (46)	372	0.8(10)
Citizens	340	2.4 (2)	58.5 (74)	245	20.4 (65)	57.6 (83)	72	* *	129	36.4(25)	558	5.2 (90)	558	5.2(71)

- (1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators ^(2, 3)														
Clinic	Prepregnancy BMI ⁽⁴⁾			Weight Gain ⁽⁵⁾			Anemia (Low Hb/Hct) ⁽⁶⁾				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Civic Ctr	868	5.3 (52)	57.3 (65)	757	18.4 (39)	58.9 (87)	175	33.7 (7)	421	21.1 (8)	917	2.8 (41)	917	3.2(49)
Clark Co	577	5.7 (60)	58.8 (78)	557	18.9 (47)	55.3 (66)	110	35.5 (8)	112	33.9(22)	704	4.4 (77)	704	1.6(23)
Clay City	5	* *	* *	5	* *	* *	0	* *	1	* *	5	* *	5	* *
Clinton	73	* *	* *	71	* *	* *	13	* *	7	* *	87	* *	87	* *
Clinton Co	220	3.2 (8)	63.2 (91)	218	17.4 (28)	49.5 (15)	51	* *	38	* *	267	3.4 (56)	267	3.0(45)
Comm East	77	* *	* *	75	* *	* *	15	* *	0	* *	118	4.2 (71)	118	13.6(103)
Cottage Gr	644	5.0 (42)	57.6 (67)	593	21.6 (76)	51.3 (30)	130	36.2 (9)	140	25.0(14)	818	3.7 (61)	818	6.4(83)
Crawford	64	* *	* *	62	* *	* *	13	* *	13	* *	80	* *	80	* *
Crown Pt	403	3.5 (15)	52.1 (23)	382	11.5 (1)	61.3 (91)	95	* *	120	13.3 (1)	538	5.6 (94)	538	4.8(67)
Daviess Co	164	2.4 (2)	62.8 (90)	153	17.0 (23)	52.9 (42)	32	* *	111	40.5(29)	187	2.1 (30)	187	5.3(75)
Dearborn	249	6.0 (68)	51.0 (19)	238	16.8 (17)	61.8 (93)	46	* *	72	* *	321	1.6 (22)	321	1.6(23)
Decatur	135	5.9 (66)	54.1 (39)	130	19.2 (52)	54.6 (62)	30	* *	27	* *	157	0.6 (7)	157	3.8(57)
Decatur Co	151	6.0 (68)	53.6 (32)	147	15.0 (7)	50.3 (23)	25	* *	32	* *	190	3.2 (51)	190	6.8(85)
Dekalb Co	253	6.7 (80)	51.4 (20)	235	21.3 (74)	56.2 (72)	40	* *	42	* *	295	1.7 (24)	295	1.4(20)
Dubois Co	191	5.8 (62)	60.7 (86)	185	16.8 (17)	51.9 (34)	41	* *	36	* *	249	4.4 (77)	249	2.8(43)
E Chicago	444	2.7 (6)	63.5 (93)	428	22.4 (83)	51.4 (31)	86	* *	57	* *	558	3.4 (56)	558	5.2(71)
Eaton	8	* *	* *	8	* *	* *	1	* *	0	* *	11	* *	11	* *

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
- (4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
- (5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
- (6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators ^(2, 3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Elkhart	1,137	4.2 (24)	58.0 (70)	1,101	22.0 (77)	50.6 (25)	125	28.8 (3)	169	16.0 (2)	1,290	2.2 (31)	1,290	2.6(38)
Elwood	71	* *	* *	70	* *	* *	17	* *	17	* *	90	* *	90	* *
Fayette Co	146	6.8 (82)	55.5 (51)	140	18.6 (43)	50.7 (27)	22	* *	33	* *	183	1.1 (11)	183	1.1(15)
Floyd Co	377	5.0 (42)	53.6 (32)	361	20.2 (62)	52.9 (42)	74	* *	77	* *	461	6.3 (98)	461	9.8(99)
Forest Man	537	3.5 (15)	61.8 (88)	497	22.1 (78)	53.7 (52)	103	33.0 (6)	102	26.5(15)	681	2.5 (35)	681	3.1(47)
Fountain	92	* *	* *	87	* *	* *	14	* *	23	* *	106	1.9 (26)	106	7.5(91)
Franklin C	81	* *	* *	80	* *	* *	12	* *	14	* *	99	* *	99	* *
Fulton Co	117	5.1 (47)	60.7 (86)	104	16.3 (15)	61.5 (92)	16	* *	36	* *	131	0.0 (1)	131	0.8(10)
Georgetown	1,000	4.0 (22)	59.3 (79)	935	19.3 (53)	51.1 (28)	241	36.5 (11)	218	23.4(11)	1,342	4.3 (74)	1,342	4.5(64)
Gibson Co	150	5.3 (52)	50.0 (11)	137	22.6 (87)	53.3 (49)	35	* *	31	* *	187	1.1 (11)	187	1.1(15)
Glenn Pk	351	4.8 (37)	58.7 (77)	247	22.3 (81)	52.2 (37)	94	* *	77	* *	490	1.2 (15)	490	3.1(47)
Goshen	519	3.7 (19)	56.6 (62)	498	15.5 (8)	52.6 (40)	66	* *	78	* *	603	3.8 (63)	603	0.8(10)
Grant Co	450	5.3 (52)	49.8 (10)	422	19.0 (50)	51.7 (32)	87	* *	238	24.8(13)	541	3.3 (54)	541	7.2(87)
Grant St	313	5.1 (47)	64.5 (94)	296	17.6 (32)	62.5 (94)	60	* *	39	* *	407	1.2 (15)	407	3.7(53)
Grassy Crk	716	4.5 (33)	60.2 (83)	682	18.9 (47)	51.9 (34)	129	36.4 (10)	153	26.8(16)	935	2.9 (43)	935	3.5(52)
Greene Co	129	6.2 (74)	54.3 (41)	123	20.3 (63)	56.1 (71)	26	* *	44	* *	161	1.9 (26)	161	0.6(7)
Hamilton	527	4.9 (39)	46.3 (2)	505	18.4 (39)	53.9 (56)	130	37.7 (13)	158	27.2(18)	663	5.1 (87)	663	3.2(49)

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
- (4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
- (5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
- (6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators ^(2, 3)														
Clinic	Prepregnancy BMI ⁽⁴⁾			Weight Gain ⁽⁵⁾			Anemia (Low Hb/Hct) ⁽⁶⁾				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight	Over-weight		< Ideal	> Ideal	3rd Trimester		Postpartum						
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Hammond	507	3.9 (20)	60.2 (83)	481	18.1 (35)	55.7 (70)	114	37.7 (13)	117	39.3(28)	647	3.2 (51)	647	2.6(38)
Hancock Co	226	4.9 (39)	53.1 (27)	216	15.7 (10)	53.2 (46)	54	* *	68	* *	288	3.1 (49)	288	1.0(14)
Hanna	1,039	5.5 (55)	51.4 (20)	987	22.2 (79)	46.9 (4)	203	26.6 (1)	260	26.9(17)	1,351	5.1 (87)	1,351	3.7(53)
Harrison	200	5.5 (55)	50.0 (11)	190	21.1 (72)	50.0 (21)	26	* *	37	* *	245	6.5 (99)	245	5.7(78)
Hendricks	310	3.2 (8)	52.9 (26)	261	11.5 (1)	58.2 (86)	88	* *	217	47.9(30)	373	2.7 (40)	373	1.6(23)
Henry Co	253	7.1 (87)	48.2 (4)	249	16.9 (20)	56.6 (77)	39	* *	26	* *	297	3.0 (46)	297	0.3(3)
Hessville	281	4.6 (34)	57.7 (69)	274	19.3 (53)	54.7 (63)	75	* *	52	* *	356	1.1 (11)	356	0.6(7)
Hispanic C	106	1.9 (1)	60.4 (85)	102	27.5 (93)	38.2 (2)	22	* *	9	* *	130	4.6 (81)	130	4.6(65)
Howard Co	570	5.6 (57)	55.3 (50)	553	17.2 (26)	50.6 (25)	82	* *	123	23.6(12)	690	1.2 (15)	690	1.3(19)
Huntington	196	7.1 (87)	49.0 (7)	187	20.9 (71)	48.7 (13)	25	* *	27	* *	224	4.5 (80)	224	5.8(80)
Jackson Co	288	5.9 (66)	55.9 (55)	276	18.8 (45)	53.3 (49)	49	* *	75	* *	344	3.8 (63)	344	5.2(71)
Jasonville	6	* *	* *	6	* *	* *	1	* *	3	* *	7	* *	7	* *
Jasper Co	171	4.1 (23)	54.4 (43)	162	17.3 (27)	56.2 (72)	14	* *	38	* *	209	0.0 (1)	209	0.0(1)
Jay Co	113	6.2 (74)	58.4 (73)	111	25.2 (91)	50.5 (24)	22	* *	38	* *	157	2.5 (35)	157	2.5(37)
Jefferson	171	7.0 (85)	55.0 (49)	162	18.5 (42)	51.9 (34)	27	* *	36	* *	219	4.1 (68)	219	5.5(76)
Jennings	243	5.8 (62)	54.7 (45)	229	18.3 (38)	53.7 (52)	21	* *	48	* *	274	2.6 (38)	274	2.2(33)
Johnson Co	526	6.8 (82)	50.8 (16)	506	15.6 (9)	56.9 (78)	133	42.1 (17)	144	19.4 (5)	712	3.8 (63)	712	8.0(94)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators ^(2, 3)														
Clinic	Prepregnancy BMI ⁽⁴⁾			Weight Gain ⁽⁵⁾			Anemia (Low Hb/Hct) ⁽⁶⁾				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Kosciusko	342	5.6 (57)	53.5 (30)	325	20.6 (68)	48.3 (10)	75	* *	106	16.0 (2)	425	4.2 (71)	425	2.8(43)
Lagrange	111	7.2 (89)	51.4 (20)	106	25.5 (92)	50.0 (21)	26	* *	35	* *	139	4.3 (74)	139	4.3(62)
Lake Station	305	6.2 (74)	59.7 (80)	235	17.0 (23)	57.0 (79)	58	* *	62	* *	444	4.3 (74)	444	7.9(93)
Laporte	344	6.1 (73)	54.1 (39)	324	17.0 (23)	53.4 (51)	75	* *	96	* *	407	3.4 (56)	407	1.7(28)
Main St	662	5.6 (57)	56.0 (58)	632	19.3 (53)	56.2 (72)	115	40.0 (16)	113	38.1(27)	816	2.8 (41)	816	3.3(51)
Marshall	296	2.4 (2)	60.1 (82)	279	15.8 (12)	54.5 (60)	60	* *	63	* *	362	4.7 (82)	362	5.5(76)
Martin Co	47	* *	* *	46	* *	* *	4	* *	1	* *	56	* *	56	* *
Memorial	147	9.5 (94)	47.6 (3)	145	22.8 (88)	38.6 (3)	28	* *	0	* *	182	1.1 (11)	182	0.5(4)
Merrillville	0	* *	* *	0	* *	* *	0	* *	0	* *	2	* *	2	* *
Methodist	141	4.3 (27)	53.9 (37)	137	19.0 (50)	59.1 (88)	27	* *	0	* *	222	6.8(100)	222	8.6(98)
Miami Co	166	6.0 (68)	50.0 (11)	160	17.5 (30)	55.0 (65)	28	* *	115	28.7(19)	197	2.5 (35)	197	6.1(81)
Mich City	352	5.7 (60)	58.2 (72)	326	16.9 (20)	54.0 (57)	72	* *	71	* *	432	1.4 (19)	432	1.6(23)
Moms/Kids	99	* *	* *	94	* *	* *	27	* *	13	* *	120	4.2 (71)	120	7.5(91)
Monroe Co	526	4.9 (39)	53.8 (35)	484	18.4 (39)	53.7 (52)	103	27.2 (2)	145	32.4(21)	615	3.3 (54)	615	0.8(10)
Montgomery	225	4.4 (30)	53.3 (28)	214	16.8 (17)	57.9 (84)	48	* *	53	* *	270	4.1 (68)	270	1.1(15)
Morgan Co	299	6.0 (68)	58.5 (74)	279	13.6 (4)	58.1 (85)	60	* *	65	* *	407	2.0 (28)	407	1.7(28)
N Arlingto	615	4.2 (24)	62.1 (89)	584	18.0 (34)	55.3 (66)	114	45.6 (20)	117	22.2 (9)	813	4.1 (68)	813	6.4(83)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators ^(2, 3)														
Clinic	Prepregnancy BMI ⁽⁴⁾			Weight Gain ⁽⁵⁾			Anemia (Low Hb/Hct) ⁽⁶⁾				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
N Liberty	16	* *	* *	16	* *	* *	1	* *	5	* *	18	* *	18	* *
N Manchest	40	* *	* *	38	* *	* *	10	* *	27	* *	50	* *	50	* *
Newton Co	43	* *	* *	43	* *	* *	7	* *	10	* *	57	* *	57	* *
Northwest	425	2.6 (5)	58.1 (71)	400	20.5 (66)	49.3 (14)	108	37.0 (12)	86	* *	601	4.7 (82)	601	4.3(62)
Ohio Co	4	* *	* *	4	* *	* *	1	* *	0	* *	7	* *	7	* *
Orange Co	122	3.3 (12)	65.6 (95)	116	22.4 (83)	47.4 (8)	16	* *	27	* *	155	5.2 (90)	155	11.6(102)
Owen Co	99	* *	* *	96	* *	* *	9	* *	25	* *	113	10.6(101)	113	11.5(101)
Parke Co	59	* *	* *	58	* *	* *	8	* *	19	* *	75	* *	75	* *
Perry Co	97	* *	* *	93	* *	* *	16	* *	22	* *	124	3.2 (51)	124	7.3(88)
Pike Co	61	* *	* *	56	* *	* *	10	* *	6	* *	69	* *	69	* *
Portage	315	4.4 (30)	52.7 (24)	293	18.8 (45)	55.3 (66)	58	* *	81	* *	382	2.6 (38)	382	3.7(53)
Posey Co	75	* *	* *	70	* *	* *	15	* *	18	* *	93	* *	93	* *
Pulaski Co	54	* *	* *	52	* *	* *	7	* *	7	* *	68	* *	68	* *
Purdue	114	7.0 (85)	25.4 (1)	107	20.6 (68)	32.7 (1)	17	* *	22	* *	127	1.6 (22)	127	0.0(1)
Putnam Co	188	3.2 (8)	48.4 (5)	184	17.4 (28)	57.1 (81)	20	* *	47	* *	219	11.0(102)	219	7.3(88)
Randolph	158	8.2 (92)	55.7 (54)	148	27.7 (94)	47.3 (6)	34	* *	30	* *	186	0.5 (5)	186	1.6(23)
Richmond	469	5.8 (62)	55.9 (55)	442	22.4 (83)	49.8 (18)	65	* *	82	* *	558	6.1 (97)	558	3.8(57)

- (1) Reporting period is January 1 through December 31.
 - (2) Excludes records with unknown data or errors.
 - (3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
 - (4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
 - (5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
 - (6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators ^(2, 3)														
Clinic	Prepregnancy BMI ⁽⁴⁾			Weight Gain ⁽⁵⁾			Anemia (Low Hb/Hct) ⁽⁶⁾				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Ripley Co	145	3.4 (13)	55.9 (55)	137	12.4 (3)	52.6 (40)	22	* *	41	* *	188	5.9 (95)	188	6.9(86)
Rush Co	89	* *	* *	88	* *	* *	13	* *	16	* *	105	4.8 (85)	105	3.8(57)
SB Ireland Rd	567	3.5 (15)	56.3 (60)	531	16.0 (14)	57.1 (81)	109	31.2 (4)	159	32.1(20)	679	2.2 (31)	679	2.7(41)
SB Lafayette	795	5.0 (42)	56.6 (62)	761	22.9 (89)	47.3 (6)	145	44.1 (19)	253	34.0(23)	969	1.5 (20)	969	2.3(36)
Scott Co	191	9.9 (95)	53.9 (37)	189	22.2 (79)	54.5 (60)	14	* *	25	* *	225	3.6 (59)	225	4.0(61)
Shelby Co	256	6.3 (77)	50.0 (11)	247	18.2 (36)	48.6 (12)	48	* *	90	* *	317	3.8 (63)	317	1.9(32)
Shelters	4	* *	* *	3	* *	* *	0	* *	0	* *	3	* *	3	* *
SJ Reg Ctr	36	* *	* *	36	* *	* *	7	* *	0	* *	47	* *	47	* *
So Calhoun	1,965	4.4 (30)	54.3 (41)	1,876	20.7 (70)	49.7 (17)	331	42.3 (18)	316	34.8(24)	2,339	2.2 (31)	2,339	3.8(57)
Spencer Co	82	* *	* *	80	* *	* *	21	* *	12	* *	102	2.9 (43)	102	4.9(69)
St Francis	0	* *	* *	0	* *	* *	0	* *	0	* *	1	* *	1	* *
St V-FLC	98	* *	* *	92	* *	* *	26	* *	2	* *	138	3.6 (59)	138	2.2(33)
Starke Co	150	6.7 (80)	49.3 (9)	146	19.9 (60)	56.2 (72)	20	* *	15	* *	167	3.0 (46)	167	1.2(18)
Steuben Co	234	5.1 (47)	56.4 (61)	227	19.8 (59)	53.7 (52)	26	* *	31	* *	269	0.7 (8)	269	0.7(9)
Sullivan	139	5.8 (62)	63.3 (92)	133	20.3 (63)	51.1 (28)	20	* *	53	* *	153	0.7 (8)	153	2.6(38)
Switz""Land	30	* *	* *	29	* *	* *	5	* *	13	* *	42	* *	42	* *
Tippecanoe	931	4.3 (27)	55.5 (51)	877	20.1 (61)	49.8 (18)	155	39.4 (15)	134	20.1 (6)	1,100	2.9 (43)	1,100	1.5(21)

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
- (4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
- (5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
- (6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators ^(2, 3)														
Clinic	Prepregnancy BMI ⁽⁴⁾				Weight Gain ⁽⁵⁾				Anemia (Low Hb/Hct) ⁽⁶⁾				Gestational Diabetes During Pregnancy	Hypertension During Pregnancy
	Under-weight		Over-weight		< Ideal		> Ideal		3rd Trimester		Postpartum			
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)		
Tipton Co	46	* *	* *	42	* *	* *	9	* *	8	* *	57	* *	57	* *
Union Co	35	* *	* *	34	* *	* *	4	* *	9	* *	47	* *	47	* *
United Fai	276	4.7 (35)	54.7 (45)	267	24.7 (90)	49.8 (18)	46	* *	70	* *	336	5.1 (87)	336	5.1(70)
Valpo	283	3.5 (15)	50.9 (18)	269	17.5 (30)	52.4 (39)	55	* *	71	* *	369	2.4 (34)	369	2.2(33)
Vigo	557	5.2 (51)	50.8 (16)	475	15.8 (12)	59.2 (89)	102	32.4 (5)	340	22.4(10)	600	1.8 (25)	600	3.7(53)
Vincennes	168	7.7 (91)	53.6 (32)	146	14.4 (5)	59.6 (90)	24	* *	108	16.7 (4)	189	4.8 (85)	189	4.8(67)
W 21st Ave	284	4.2 (24)	53.5 (30)	268	20.5 (66)	49.6 (16)	73	* *	27	* *	339	0.9 (10)	339	4.7(66)
Wabash	142	6.3 (77)	54.9 (47)	134	15.7 (10)	52.2 (37)	23	* *	101	20.8 (7)	158	4.4 (77)	158	8.2(96)
Warren Co	50	* *	* *	50	* *	* *	7	* *	21	* *	67	* *	67	* *
Warrick Co	205	3.4 (13)	52.7 (24)	193	17.6 (32)	56.5 (76)	36	* *	32	* *	236	0.4 (4)	236	3.0(45)
Washington	145	4.8 (37)	58.6 (76)	138	19.6 (57)	47.1 (5)	14	* *	20	* *	163	3.7 (61)	163	9.8(99)
Wells Co	119	5.0 (42)	54.6 (44)	114	21.1 (72)	55.3 (66)	30	* *	26	* *	155	1.3 (18)	155	5.2(71)
Westside	445	4.7 (35)	59.8 (81)	400	19.5 (56)	51.8 (33)	81	* *	88	* *	665	6.0 (96)	665	6.2(82)
White Co	177	2.8 (7)	49.2 (8)	173	21.4 (75)	53.2 (46)	17	* *	26	* *	208	0.5 (5)	208	0.5(4)
Whitley Co	150	6.0 (68)	53.3 (28)	142	16.9 (20)	57.0 (79)	28	* *	47	* *	178	3.9 (67)	178	1.7(28)
Wishard	85	* *	* *	81	* *	* *	22	* *	0	* *	123	11.4(103)	123	5.7(78)
Women""s	1	* *	* *	1	* *	* *	0	* *	0	* *	1	* *	1	* *

- (1) Reporting period is January 1 through December 31.
 - (2) Excludes records with unknown data or errors.
 - (3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
 - (4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
 - (5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
 - (6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators ^(2, 3)																	
Clinic	Prepregnancy BMI (4)			Weight Gain (5)				Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy			
	Under-weight		Over-weight	< Ideal		> Ideal		3rd Trimester		Postpartum							
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	
Unknown Clinics	295	8.8 (93)	48.5 (6)	1	*	*	*	*	78	*	*	2	*	*	3	*	*
Indiana	35,909	5.0	55.4	33,494	19.1	52.8	6,722	36.2	8,527	26.8	44,078	3.3	44,078	3.7			
Nation (Prior Year)	1,106,370	4.5	53.4	835,927	21.5	48.0	164,201	33.9	346,981	28.1	419,956	5.2	467,675	6.3			

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6A
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators ^(2, 3)										
Clinic	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Action Ctr	17	* *	* *	* *	31	* *	31	* *	31	* *
Albion	178	64.6 (46)	25.3 (34)		248	44.4 (44)	250	18.4 (29)	250	88.8 (44)
Anderson	515	65.0 (45)	23.3 (26)		667	41.1 (58)	666	15.5 (55)	666	90.1 (34)
Bartholome	354	44.6 (77)	45.2 (74)		425	44.5 (43)	425	18.6 (27)	426	90.1 (34)
Bedford	177	59.3 (56)	36.2 (58)		234	54.7 (10)	237	10.1 (86)	238	81.9 (85)
Benton Co	26	* *	* *	* *	35	* *	36	* *	36	* *
Berne	28	* *	* *	* *	33	* *	32	* *	32	* *
Bicknell	17	* *	* *	* *	26	* *	26	* *	26	* *
Blackburn	235	69.8 (25)	21.3 (21)		341	37.8 (74)	339	13.6 (66)	339	87.0 (64)
Blackford	63	* *	* *	* *	81	* *	81	* *	81	* *
Boone Co	122	58.2 (60)	37.7 (62)		159	49.1 (23)	159	6.3 (94)	159	90.6 (27)
Brazil	134	52.2 (70)	41.8 (68)		196	53.6 (13)	195	17.9 (33)	196	89.8 (36)
Brown Co	42	* *	* *	* *	50	* *	51	* *	51	* *
Cambridge	15	* *	* *	* *	20	* *	20	* *	20	* *
Carroll Co	50	* *	* *	* *	68	* *	66	* *	66	* *
Cass Co	256	58.6 (58)	34.0 (55)		327	45.0 (37)	326	11.0 (82)	326	82.8 (81)
Citizens	208	65.9 (39)	26.9 (40)		346	32.7 (90)	343	15.5 (55)	343	89.5 (38)
Civic Ctr	577	71.9 (19)	20.8 (17)		883	36.4 (79)	869	13.7 (65)	868	90.9 (25)
Clark Co	431	65.7 (41)	29.5 (46)		581	41.3 (54)	578	19.6 (21)	578	92.0 (21)
Clay City	4	* *	* *	* *	5	* *	5	* *	5	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6A
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators ^(2, 3)										
Clinic	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Clinton	47	* *	* *	* *	73	* *	73	* *	73	* *
Clinton Co	168	63.1 (51)	29.8 (48)		220	38.6 (69)	221	20.8 (15)	221	92.8 (15)
Comm East	61	* *	* *	* *	77	* *	77	* *	77	* *
Cottage Gr	478	70.3 (24)	21.1 (20)		638	37.5 (75)	643	7.6 (92)	642	89.3 (40)
Crawford	50	* *	* *	* *	65	* *	64	* *	64	* *
Crown Pt	290	76.6 (5)	15.5 (5)		406	33.7 (86)	402	30.1 (1)	403	95.0 (3)
Daviess Co	110	77.3 (4)	17.3 (8)		164	46.3 (33)	164	12.2 (73)	164	81.1 (89)
Dearborn	144	75.0 (8)	20.1 (13)		252	36.5 (78)	249	18.1 (31)	249	90.4 (29)
Decatur	104	53.8 (68)	43.3 (71)		137	48.9 (25)	135	23.7 (7)	135	94.8 (5)
Decatur Co	119	66.4 (37)	30.3 (49)		156	42.3 (50)	150	14.0 (62)	151	84.1 (77)
Dekalb Co	190	67.4 (35)	28.4 (43)		250	50.4 (21)	255	17.6 (36)	254	89.8 (36)
Dubois Co	124	75.0 (8)	21.0 (19)		193	34.2 (84)	191	22.5 (9)	191	86.9 (65)
E Chicago	319	68.7 (31)	24.5 (30)		447	34.9 (81)	310	16.5 (44)	428	87.6 (60)
Eaton	3	* *	* *	* *	8	* *	8	* *	8	* *
Elkhart	974	38.0 (79)	58.4 (81)		1,135	59.1 (4)	1,139	25.4 (3)	1,139	81.5 (87)
Elwood	48	* *	* *	* *	72	* *	71	* *	71	* *
Fayette Co	90	* *	* *	* *	145	51.7 (17)	146	11.6 (77)	146	86.3 (69)
Floyd Co	267	65.9 (39)	25.8 (37)		385	40.3 (61)	378	20.4 (18)	378	89.2 (41)
Forest Man	378	76.2 (6)	16.9 (6)		541	37.9 (71)	538	7.2 (93)	539	88.5 (49)
Fountain	80	* *	* *	* *	93	* *	92	* *	92	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6A
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators ^(2, 3)										
Clinic	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Franklin C	58	* *	* *	* *	82	* *	81	* *	81	* *
Fulton Co	74	* *	* *	* *	120	46.7 (31)	117	15.4 (57)	117	88.0 (54)
Georgetown	739	70.8 (21)	20.7 (16)		999	30.0 (92)	996	11.1 (81)	1,000	88.7 (45)
Gibson Co	94	* *	* *	* *	154	39.6 (65)	151	20.5 (17)	151	94.7 (6)
Glenn Pk	176	73.3 (17)	17.6 (9)		349	32.7 (90)	351	9.7 (87)	352	87.8 (59)
Goshen	450	44.9 (76)	51.1 (76)		519	53.4 (14)	519	12.1 (75)	519	81.1 (89)
Grant Co	304	68.8 (30)	23.7 (27)		451	41.2 (56)	451	16.4 (47)	451	88.5 (49)
Grant St	192	74.5 (10)	13.0 (2)		319	34.8 (82)	313	8.9 (89)	314	85.0 (75)
Grassy Crk	546	66.1 (38)	27.8 (41)		718	37.9 (71)	715	11.5 (78)	718	86.9 (65)
Greene Co	71	* *	* *	* *	131	45.0 (37)	100	11.0 (82)	127	93.7 (12)
Hamilton	362	74.0 (14)	21.3 (21)		533	37.5 (75)	526	21.5 (12)	528	92.2 (18)
Hammond	346	69.4 (27)	20.5 (15)		511	34.4 (83)	505	14.1 (61)	507	88.0 (54)
Hancock Co	181	64.1 (49)	31.5 (51)		226	42.5 (49)	228	20.6 (16)	228	90.8 (26)
Hanna	853	55.1 (65)	36.6 (60)		1,038	39.8 (63)	1,029	12.6 (71)	1,041	82.8 (81)
Harrison	137	50.4 (73)	38.7 (64)		202	47.5 (29)	201	24.9 (5)	201	96.0 (2)
Hendricks	194	82.0 (2)	11.3 (1)		314	29.0 (93)	306	22.5 (9)	311	92.6 (17)
Henry Co	215	61.9 (54)	36.3 (59)		254	52.4 (15)	254	15.0 (58)	254	81.5 (87)
Hessville	206	68.0 (33)	22.8 (25)		289	33.9 (85)	281	17.4 (39)	282	94.0 (11)
Hispanic C	86	* *	* *	* *	105	38.1 (70)	107	19.6 (21)	107	93.5 (14)
Howard Co	423	62.4 (53)	33.1 (54)		574	46.0 (34)	571	9.6 (88)	572	87.9 (56)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6A
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators ^(2, 3)									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy ⁽⁴⁾		During Pregnancy ⁽⁵⁾	
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Huntington	149	44.3 (78)	54.4 (79)	195	61.5 (3)	196	21.4 (13)	197	85.3 (74)
Jackson Co	212	74.5 (10)	20.8 (17)	291	44.7 (40)	288	13.2 (69)	289	92.0 (21)
Jasonville	3	* *	* *	6	* *	6	* *	6	* *
Jasper Co	121	62.8 (52)	35.5 (57)	170	48.2 (27)	171	14.0 (62)	171	85.4 (73)
Jay Co	83	* *	* *	112	49.1 (23)	113	11.5 (78)	113	89.4 (39)
Jefferson	141	51.8 (71)	44.0 (72)	176	54.0 (12)	172	16.9 (41)	172	83.7 (78)
Jennings	195	29.2 (82)	67.7 (82)	239	67.4 (1)	243	16.0 (49)	243	74.9 (94)
Johnson Co	418	63.4 (50)	29.4 (45)	532	33.3 (88)	508	16.5 (44)	526	90.3 (30)
Kosciusko	248	76.2 (6)	16.9 (6)	340	45.0 (37)	343	15.7 (53)	342	86.5 (67)
Lagrange	76	* *	* *	114	43.9 (46)	111	18.9 (25)	111	86.5 (67)
Lake Station	172	65.7 (41)	26.7 (39)	311	39.5 (66)	306	13.4 (68)	306	87.6 (60)
Laporte	288	54.2 (66)	40.3 (66)	345	44.6 (41)	342	18.7 (26)	343	95.0 (3)
Main St	459	74.5 (10)	21.4 (23)	665	47.4 (30)	659	17.8 (35)	663	90.2 (32)
Marshall	241	60.6 (55)	35.3 (56)	296	40.2 (62)	296	19.3 (23)	297	88.2 (53)
Martin Co	32	* *	* *	45	* *	47	* *	47	* *
Memorial	112	64.3 (48)	28.6 (44)	150	38.7 (68)	147	16.3 (48)	147	94.6 (7)
Merrillville	0	* *	* *	0	* *	0	* *	0	* *
Methodist	106	65.1 (44)	24.5 (30)	138	40.6 (60)	141	10.6 (85)	141	83.0 (80)
Miami Co	111	67.6 (34)	30.6 (50)	168	51.2 (18)	166	17.5 (37)	166	89.2 (41)
Mich City	264	58.3 (59)	33.0 (53)	352	37.2 (77)	323	21.7 (11)	345	92.8 (15)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6A
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators ^(2, 3)										
Clinic	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Moms/Kids	70	* *	* *	* *	99	* *	99	* *	99	* *
Monroe Co	349	69.6 (26)	25.8 (37)		522	44.4 (44)	380	23.4 (8)	508	92.1 (19)
Montgomery	185	52.4 (69)	43.2 (70)		222	44.6 (41)	225	16.0 (49)	225	90.2 (32)
Morgan Co	224	69.2 (29)	24.1 (28)		303	35.0 (80)	300	21.0 (14)	300	90.3 (30)
N Arlingto	475	74.3 (13)	20.4 (14)		614	41.2 (56)	583	12.7 (70)	619	87.1 (62)
N Liberty	14	* *	* *	* *	16	* *	16	* *	16	* *
N Manchest	30	* *	* *	* *	41	* *	40	* *	40	* *
Newton Co	33	* *	* *	* *	40	* *	43	* *	43	* *
Northwest	325	72.3 (18)	18.8 (11)		430	28.4 (94)	429	14.9 (60)	429	92.1 (19)
Ohio Co	4	* *	* *	* *	4	* *	4	* *	4	* *
Orange Co	88	* *	* *	* *	120	54.2 (11)	121	16.5 (44)	122	82.8 (81)
Owen Co	73	* *	* *	* *	98	* *	99	* *	99	* *
Parke Co	40	* *	* *	* *	59	* *	59	* *	58	* *
Perry Co	83	* *	* *	* *	99	* *	98	* *	98	* *
Pike Co	36	* *	* *	* *	62	* *	61	* *	61	* *
Portage	210	65.2 (43)	29.5 (46)		320	45.9 (35)	315	17.5 (37)	315	87.9 (56)
Posey Co	56	* *	* *	* *	76	* *	75	* *	75	* *
Pulaski Co	32	* *	* *	* *	54	* *	54	* *	54	* *
Purdue	86	* *	* *	* *	112	56.3 (7)	79	* *	105	97.1 (1)
Putnam Co	144	47.9 (74)	48.6 (75)		187	52.4 (15)	188	25.0 (4)	187	83.4 (79)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6A
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators ^(2, 3)										
Clinic	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Prepregnancy ⁽⁴⁾		During Pregnancy ⁽⁵⁾	
	Number	% (Rank)	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Randolph	106	82.1 (1)	13.2 (3)		157	42.7 (47)	157	10.8 (84)	158	93.7 (12)
Richmond	400	38.0 (79)	52.3 (78)		467	55.0 (8)	405	13.6 (66)	451	73.8 (95)
Ripley Co	106	67.0 (36)	28.3 (42)		146	39.0 (67)	147	18.4 (29)	147	82.3 (84)
Rush Co	63	* *	* *		86	* *	88	* *	89	* *
SB Ireland Rd	442	71.9 (19)	24.4 (29)		572	37.9 (71)	567	19.0 (24)	567	94.5 (8)
SB Lafayette	617	68.6 (32)	24.6 (32)		803	42.2 (51)	797	15.8 (52)	797	90.6 (27)
Scott Co	169	56.8 (63)	40.8 (67)		192	58.9 (5)	192	15.6 (54)	192	81.8 (86)
Shelby Co	170	69.4 (27)	24.7 (33)		252	41.3 (54)	257	8.9 (89)	256	85.9 (72)
Shelters	2	* *	* *		4	* *	4	* *	4	* *
SJ Reg Ctr	29	* *	* *		36	* *	36	* *	36	* *
So Calhoun	1,495	57.8 (62)	37.5 (61)		1,967	46.4 (32)	1,966	16.7 (43)	1,967	88.5 (49)
Spencer Co	66	* *	* *		81	* *	82	* *	82	* *
St Francis	0	* *	* *		0	* *	0	* *	0	* *
St V-FLC	72	* *	* *		98	* *	98	* *	97	* *
Starke Co	126	54.0 (67)	44.4 (73)		149	55.0 (8)	150	16.0 (49)	150	79.3 (92)
Steuben Co	177	46.3 (75)	52.0 (77)		234	62.4 (2)	234	15.0 (58)	234	79.9 (91)
Sullivan	74	* *	* *		138	50.7 (20)	138	18.1 (31)	139	87.1 (62)
Switz""Land	25	* *	* *		31	* *	31	* *	31	* *
Tippecanoe	747	50.5 (72)	43.1 (69)		925	51.1 (19)	903	19.8 (20)	929	84.7 (76)
Tipton Co	26	* *	* *		45	* *	46	* *	46	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6A
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators ^(2, 3)										
Clinic	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Union Co	22	* *	* *	* *	34	* *	35	* *	35	* *
United Fai	220	64.5 (47)	32.3 (52)		274	42.7 (47)	276	20.3 (19)	276	94.2 (9)
Valpo	206	70.4 (23)	25.7 (36)		279	39.8 (63)	283	24.7 (6)	283	91.2 (24)
Vigo	312	73.4 (16)	18.6 (10)		559	41.5 (53)	559	11.3 (80)	560	88.6 (46)
Vincennes	107	78.5 (3)	19.6 (12)		174	42.0 (52)	171	8.8 (91)	171	88.9 (43)
W 21st Ave	190	70.5 (22)	15.3 (4)		281	33.5 (87)	285	12.3 (72)	284	78.9 (93)
Wabash	97	* *	* *	* *	142	47.9 (28)	142	16.9 (41)	142	91.5 (23)
Warren Co	44	* *	* *	* *	50	* *	50	* *	50	* *
Warrick Co	141	56.7 (64)	38.3 (63)		206	45.6 (36)	205	12.2 (73)	206	87.9 (56)
Washington	118	35.6 (81)	55.1 (80)		140	57.9 (6)	145	17.9 (33)	145	86.2 (70)
Wells Co	90	* *	* *	* *	122	41.0 (59)	119	18.5 (28)	120	94.2 (9)
Westside	322	58.7 (57)	25.5 (35)		444	32.9 (89)	446	13.9 (64)	446	88.6 (46)
White Co	126	73.8 (15)	22.2 (24)		176	48.3 (26)	178	11.8 (76)	178	86.0 (71)
Whitley Co	112	58.0 (61)	40.2 (65)		149	50.3 (22)	150	28.0 (2)	149	88.6 (46)
Wishard	72	* *	* *	* *	85	* *	85	* *	85	* *
Women""s	1	* *	* *	* *	1	* *	1	* *	1	* *
Unknown Clinics	1	* *	* *	* *	305	16.7 (95)	294	17.0 (40)	296	88.5 (49)
Indiana	26,281	62.8	30.8		36,028	42.6	35,411	16.1	35,904	88.0

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6A
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators ^(2, 3)									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Pregpregnancy (4)		During Pregpregnancy (5)	
	Number	%	%	Number	%	Number	%	Number	%
	(Rank)	(Rank)	(Rank)	(Rank)	(Rank)	(Rank)	(Rank)	(Rank)	(Rank)
Nation (Prior Year)	429,770	83.1	4.1	1,074,863	36.0	380,154	19.2	418,448	85.0

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2011 Pregnancy Nutrition Surveillance (1)
Indiana
Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Action Ctr	40	* *	36	* *	8	* *	30	* *	35	* *
Albion	304	34.5 (55)	291	27.5 (77)	99	* *	250	47.6 (58)	290	36.9 (51)
Anderson	866	35.8 (62)	841	25.7 (61)	295	32.5 (28)	666	50.6 (70)	835	44.6 (73)
Bartholome	526	37.5 (69)	516	26.6 (69)	192	30.2 (35)	426	54.7 (82)	510	50.4 (87)
Bedford	285	41.4 (90)	283	29.0 (85)	117	35.0 (18)	238	45.0 (54)	280	42.9 (64)
Benton Co	45	* *	43	* *	16	* *	36	* *	43	* *
Berne	38	* *	38	* *	9	* *	32	* *	38	* *
Bicknell	30	* *	26	* *	14	* *	26	* *	26	* *
Blackburn	454	20.0 (15)	437	12.4 (15)	88	* *	336	31.0 (24)	435	26.2 (31)
Blackford	102	38.2 (78)	99	* *	38	* *	81	* *	99	* *
Boone Co	202	41.1 (88)	198	33.3 (94)	81	* *	159	51.6 (74)	195	51.8 (92)
Brazil	231	37.7 (70)	225	25.8 (63)	84	* *	196	58.7 (91)	222	57.2 (99)
Brown Co	64	* *	62	* *	25	* *	51	* *	61	* *
Cambridge	28	* *	28	* *	8	* *	20	* *	27	* *
Carroll Co	82	* *	81	* *	23	* *	66	* *	80	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2011 Pregnancy Nutrition Surveillance (1)
Indiana
Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Cass Co	384	26.3 (31)	372	18.0 (28)	95	* *	326	31.6 (26)	366	24.9 (30)
Citizens	654	16.1 (8)	558	16.8 (23)	74	* *	340	39.1 (43)	555	35.0 (46)
Civic Ctr	1,021	39.0 (80)	917	19.3 (35)	347	45.8 (8)	868	22.4 (16)	906	7.9 (7)
Clark Co	722	31.6 (45)	704	22.4 (48)	216	33.8 (24)	578	38.9 (42)	699	35.6 (48)
Clay City	5	* *	5	* *	1	* *	5	* *	5	* *
Clinton	89	* *	87	* *	30	* *	72	* *	87	* *
Clinton Co	269	26.0 (28)	267	18.4 (30)	70	* *	221	20.8 (10)	261	6.1 (5)
Comm East	119	30.3 (42)	118	23.7 (56)	35	* *	59	* *	117	43.6 (67)
Cottage Gr	864	30.2 (41)	818	31.2 (92)	239	19.2 (46)	635	46.1 (56)	806	46.0 (75)
Crawford	82	* *	80	* *	29	* *	64	* *	78	* *
Crown Pt	558	26.9 (34)	538	12.6 (16)	144	58.3 (2)	403	40.7 (45)	529	35.3 (47)
Daviess Co	196	37.8 (72)	187	23.5 (55)	69	* *	164	48.8 (65)	184	46.2 (76)
Dearborn	330	42.7 (93)	321	29.0 (85)	136	41.2 (11)	249	48.2 (63)	312	43.6 (67)
Decatur	161	32.9 (48)	157	19.7 (37)	52	* *	135	48.9 (66)	155	43.9 (71)
Decatur Co	194	38.1 (77)	190	28.9 (84)	71	* *	151	58.9 (92)	188	55.9 (98)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2011 Pregnancy Nutrition Surveillance (1)
Indiana
Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Dekalb Co	306	33.0 (50)	295	27.5 (77)	99	* *	255	58.4 (90)	292	49.3 (86)
Dubois Co	252	27.0 (35)	249	21.3 (45)	66	* *	186	43.0 (48)	246	40.7 (60)
E Chicago	568	14.8 (7)	558	5.9 (4)	83	* *	445	21.1 (11)	549	13.1 (12)
Eaton	11	* *	11	* *	3	* *	8	* *	11	* *
Elkhart	1,321	26.1 (29)	1,290	16.3 (21)	330	29.1 (38)	1,136	21.3 (13)	1,276	9.3 (8)
Elwood	91	* *	90	* *	36	* *	71	* *	87	* *
Fayette Co	189	36.5 (66)	183	25.7 (61)	66	* *	146	61.6 (95)	182	54.4 (95)
Floyd Co	475	38.3 (79)	461	26.0 (66)	174	37.9 (15)	377	43.5 (50)	456	41.9 (62)
Forest Man	719	20.4 (18)	681	11.3 (11)	134	38.8 (14)	332	21.7 (14)	495	21.6 (25)
Fountain	111	47.7 (104)	106	34.0 (96)	52	* *	91	* *	105	60.0 (100)
Franklin C	100	38.0 (75)	99	* *	38	* *	81	* *	97	* *
Fulton Co	143	36.4 (65)	131	26.0 (66)	49	* *	117	47.9 (60)	131	42.0 (63)
Georgetown	1,402	11.8 (5)	1,342	7.3 (6)	150	50.0 (4)	996	18.6 (9)	1,328	18.4 (18)
Gibson Co	200	43.5 (95)	187	39.0 (101)	79	* *	151	57.0 (87)	186	55.4 (97)
Glenn Pk	594	10.9 (4)	490	6.9 (5)	42	* *	352	23.6 (17)	485	20.2 (22)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2011 Pregnancy Nutrition Surveillance (1)
Indiana
Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Goshen	622	20.1 (16)	603	11.4 (12)	122	40.2 (12)	518	15.1 (3)	599	5.2 (3)
Grant Co	564	33.0 (50)	541	27.0 (75)	176	26.7 (40)	451	35.5 (35)	535	9.7 (9)
Grant St	415	16.4 (9)	407	13.3 (18)	67	* *	313	26.5 (20)	401	26.9 (32)
Grassy Crk	965	18.5 (13)	935	9.7 (9)	171	49.7 (5)	714	25.9 (19)	928	22.4 (27)
Greene Co	163	39.3 (83)	161	26.1 (68)	61	* *	129	61.2 (94)	159	50.9 (88)
Hamilton	682	22.0 (21)	663	12.2 (14)	143	47.6 (6)	527	27.7 (21)	653	19.9 (21)
Hammond	669	18.4 (12)	647	8.3 (7)	117	59.8 (1)	505	22.0 (15)	637	19.2 (20)
Hancock Co	296	29.4 (40)	288	19.1 (33)	85	* *	226	24.8 (18)	286	14.0 (15)
Hanna	1,393	20.9 (19)	1,351	17.2 (26)	275	30.9 (33)	1,039	34.6 (32)	1,324	30.3 (37)
Harrison	254	43.3 (94)	245	29.8 (88)	106	34.9 (19)	201	49.8 (69)	240	34.2 (44)
Hendricks	421	27.6 (36)	373	15.3 (20)	94	* *	311	39.9 (44)	368	38.9 (55)
Henry Co	300	47.7 (104)	297	33.7 (95)	140	26.4 (41)	253	53.4 (79)	294	48.0 (79)
Hessville	364	22.5 (22)	356	9.8 (10)	78	* *	281	16.4 (5)	355	11.3 (10)
Hispanic C	133	6.0 (2)	130	0.8 (1)	8	* *	107	15.9 (4)	127	7.1 (6)
Howard Co	705	35.6 (61)	690	26.7 (70)	245	27.3 (39)	572	47.9 (60)	680	36.8 (50)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2011 Pregnancy Nutrition Surveillance (1)
Indiana
Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Huntington	232	46.1 (101)	224	28.6 (82)	101	31.7 (32)	197	52.8 (77)	221	51.6 (91)
Jackson Co	352	36.1 (63)	344	22.4 (48)	123	39.0 (13)	289	38.8 (40)	341	33.1 (41)
Jasonville	7	* *	7	* *	4	* *	6	* *	7	* *
Jasper Co	217	35.5 (60)	209	28.7 (83)	74	* *	171	60.8 (93)	208	46.6 (78)
Jay Co	158	32.9 (48)	157	26.8 (73)	51	* *	113	48.7 (64)	153	37.3 (52)
Jefferson	224	44.6 (97)	219	36.5 (99)	96	* *	172	58.1 (88)	213	43.7 (69)
Jennings	284	39.8 (86)	274	31.4 (93)	108	18.5 (47)	243	56.8 (86)	271	48.7 (82)
Johnson Co	730	27.8 (37)	712	21.2 (44)	195	32.8 (26)	527	42.7 (47)	699	29.9 (36)
Kosciusko	441	26.1 (29)	425	20.9 (41)	110	25.5 (43)	343	38.5 (38)	420	23.3 (28)
Lagrange	144	26.4 (32)	139	20.9 (41)	35	* *	111	43.2 (49)	134	39.6 (57)
Lake Station	512	20.1 (16)	444	18.0 (28)	80	* *	306	47.1 (57)	442	40.5 (59)
Laporte	423	37.1 (68)	407	27.5 (77)	146	30.8 (34)	344	51.7 (75)	398	48.0 (79)
Main St	837	33.9 (54)	816	25.5 (60)	277	23.8 (44)	662	30.7 (23)	801	14.2 (16)
Marshall	373	28.7 (38)	362	19.9 (38)	100	34.0 (22)	296	17.6 (8)	361	5.8 (4)
Martin Co	57	* *	56	* *	25	* *	47	* *	55	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2011 Pregnancy Nutrition Surveillance (1)
Indiana
Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Memorial	183	25.1 (25)	182	14.8 (19)	46	* *	147	17.0 (7)	182	13.7 (13)
Merrillville	2	* *	2	* *	0	* *	0	* *	2	* *
Methodist	225	21.8 (20)	222	23.0 (53)	49	* *	127	41.7 (46)	222	44.1 (72)
Miami Co	202	46.0 (100)	197	25.4 (59)	89	* *	166	56.6 (84)	197	46.2 (76)
Mich City	455	28.8 (39)	432	22.0 (47)	121	35.5 (17)	350	35.4 (34)	426	30.3 (37)
Moms/Kids	125	23.2 (23)	120	9.2 (8)	26	* *	99	* *	119	24.4 (29)
Monroe Co	657	33.6 (53)	615	22.9 (52)	201	31.8 (29)	526	33.7 (29)	612	22.1 (26)
Montgomery	282	37.9 (74)	270	27.0 (75)	101	34.7 (20)	224	56.7 (85)	268	43.3 (66)
Morgan Co	423	34.5 (55)	407	30.2 (89)	136	29.4 (36)	298	51.0 (71)	403	48.4 (81)
N Arlingto	838	23.7 (24)	813	18.5 (31)	189	33.9 (23)	485	34.0 (30)	794	35.6 (48)
N Liberty	18	* *	18	* *	9	* *	16	* *	18	* *
N Manchest	52	* *	50	* *	15	* *	40	* *	50	* *
Newton Co	57	* *	57	* *	12	* *	43	* *	57	* *
Northwest	624	7.7 (3)	601	4.5 (3)	43	* *	425	12.5 (2)	598	13.9 (14)
Ohio Co	7	* *	7	* *	0	* *	4	* *	7	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2011 Pregnancy Nutrition Surveillance (1)
Indiana
Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Orange Co	159	35.2 (59)	155	28.4 (81)	52	* *	122	32.8 (27)	152	19.1 (19)
Owen Co	116	50.0 (106)	113	39.8 (103)	57	* *	99	* *	113	51.3 (90)
Parke Co	75	* *	75	* *	33	* *	58	* *	75	* *
Perry Co	127	46.5 (102)	124	39.5 (102)	56	* *	98	* *	123	61.8 (102)
Pike Co	74	* *	69	* *	28	* *	61	* *	67	* *
Portage	403	33.3 (52)	382	18.6 (32)	126	50.8 (3)	313	47.6 (58)	374	40.4 (58)
Posey Co	98	* *	93	* *	41	* *	75	* *	93	* *
Pulaski Co	70	* *	68	* *	16	* *	54	* *	67	* *
Purdue	133	3.8 (1)	127	0.8 (1)	5	* *	114	4.4 (1)	126	4.0 (1)
Putnam Co	222	44.6 (97)	219	28.3 (80)	98	* *	188	51.1 (72)	217	48.8 (83)
Randolph	190	32.1 (46)	186	24.2 (57)	60	* *	158	48.1 (62)	177	37.9 (54)
Richmond	577	39.2 (81)	558	26.7 (70)	214	29.4 (36)	470	51.5 (73)	550	37.5 (53)
Ripley Co	196	39.3 (83)	188	35.1 (97)	70	* *	146	58.2 (89)	182	53.8 (94)
Rush Co	106	47.2 (103)	105	30.5 (90)	49	* *	89	* *	103	55.3 (96)
SB Ireland Rd	711	30.9 (43)	679	17.2 (26)	208	43.8 (10)	564	31.4 (25)	673	21.2 (24)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2011 Pregnancy Nutrition Surveillance (1)
Indiana
Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
SB Lafayette	999	25.2 (26)	969	13.0 (17)	238	46.2 (7)	797	21.2 (12)	948	12.7 (11)
Scott Co	227	41.9 (92)	225	35.6 (98)	93	* *	192	55.2 (83)	224	52.7 (93)
Shelby Co	324	37.0 (67)	317	25.9 (65)	113	32.7 (27)	257	52.5 (76)	313	51.1 (89)
Shelters	4	* *	3	* *	2	* *	4	* *	3	* *
SJ Reg Ctr	47	* *	47	* *	15	* *	36	* *	47	* *
So Calhoun	2,409	26.6 (33)	2,339	17.0 (24)	610	33.0 (25)	1,959	36.1 (36)	2,302	33.1 (41)
Spencer Co	102	36.3 (64)	102	22.5 (50)	37	* *	81	* *	97	* *
St Francis	1	* *	1	* *	0	* *	0	* *	1	* *
St V-FLC	142	16.9 (10)	138	16.7 (22)	23	* *	88	* *	138	32.6 (40)
Starke Co	171	35.1 (58)	167	26.9 (74)	59	* *	150	46.0 (55)	166	48.8 (83)
Steuben Co	275	39.6 (85)	269	22.7 (51)	107	31.8 (29)	233	54.1 (80)	263	45.6 (74)
Sullivan	158	44.3 (96)	153	36.6 (100)	67	* *	139	53.2 (78)	153	49.0 (85)
Switz""Land	44	* *	42	* *	17	* *	31	* *	40	* *
Tippecanoe	1,140	25.6 (27)	1,100	20.2 (39)	278	22.7 (45)	932	38.8 (40)	1,084	33.8 (43)
Tipton Co	61	* *	57	* *	17	* *	46	* *	57	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2011 Pregnancy Nutrition Surveillance (1)
Indiana
Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Union Co	48	* *	47	* *	10	* *	35	* *	46	* *
United Fai	342	32.5 (47)	336	21.1 (43)	107	26.2 (42)	276	37.3 (37)	331	32.0 (39)
Valpo	382	34.6 (57)	369	19.2 (34)	128	45.3 (9)	283	16.6 (6)	360	4.2 (2)
Vigo	676	37.7 (70)	600	23.2 (54)	218	34.4 (21)	560	49.5 (68)	592	42.9 (64)
Vincennes	209	37.8 (72)	189	24.9 (58)	73	* *	170	43.5 (50)	184	41.8 (61)
W 21st Ave	354	17.5 (11)	339	17.1 (25)	60	* *	284	34.9 (33)	338	28.1 (35)
Wabash	165	40.0 (87)	158	31.0 (91)	64	* *	142	33.1 (28)	156	20.5 (23)
Warren Co	67	* *	67	* *	29	* *	49	* *	67	* *
Warrick Co	246	45.1 (99)	236	26.7 (70)	107	31.8 (29)	206	54.4 (81)	233	61.4 (101)
Washington	168	41.1 (88)	163	25.8 (63)	66	* *	145	44.1 (53)	162	27.2 (33)
Wells Co	158	41.8 (91)	155	29.0 (85)	65	* *	120	49.2 (67)	153	43.8 (70)
Westside	709	18.5 (13)	665	19.4 (36)	118	36.4 (16)	444	34.0 (30)	655	34.5 (45)
White Co	213	31.5 (44)	208	20.7 (40)	66	* *	178	29.8 (22)	207	15.0 (17)
Whitley Co	184	38.0 (75)	178	21.3 (45)	68	* *	150	44.0 (52)	177	39.0 (56)
Wishard	124	12.9 (6)	123	11.4 (12)	15	* *	80	* *	123	27.6 (34)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2011 Pregnancy Nutrition Surveillance (1)
Indiana
Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Women""s	1	* *	1	* *	0	* *	1	* *	1	* *
Unknown Clinics	306	39.2 (81)	3	* *	1	* *	291	38.5 (38)	3	* *
Indiana	46,227	29.0	44,078	20.3	12,540	33.8	35,519	37.4	43,331	31.1
Nation (Prior Year)	814,631	23.0	715,034	12.1	137,883	38.1	660,570	16.3	562,126	15.6

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators ^(2, 3)											
Clinic	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed		
	Low (4)		High (5)		Number	%	Number	%	Number	%	
	Number	% (Rank)	% (Rank)	(Rank)							(Rank)
Action Ctr	34	* * * *	* *		25	* *		24	* *	34	* *
Albion	283	7.1 (35)	5.3 (26)		233	13.3 (55)		202	4.0 (73)	283	73.9 (26)
Anderson	821	8.5 (62)	5.6 (31)		636	14.9 (70)		534	1.9 (21)	829	59.7 (84)
Bartholome	498	6.4 (19)	6.6 (51)		408	12.3 (40)		358	2.5 (37)	498	70.9 (43)
Bedford	274	9.9 (81)	6.6 (51)		229	9.6 (11)		207	4.3 (77)	274	70.4 (45)
Benton Co	39	* * * *	* *		31	* *		23	* *	39	* *
Berne	38	* * * *	* *		33	* *		30	* *	38	* *
Bicknell	24	* * * *	* *		20	* *		19	* *	24	* *
Blackburn	432	10.2 (88)	4.4 (8)		321	14.0 (64)		276	2.9 (49)	433	46.7 (98)
Blackford	99	* * * *	* *		78	* *		71	* *	99	* *
Boone Co	189	5.3 (9)	6.9 (61)		151	13.9 (62)		130	2.3 (31)	189	56.1 (92)
Brazil	213	5.6 (14)	4.7 (13)		181	12.7 (43)		158	1.9 (21)	213	64.8 (65)
Brown Co	59	* * * *	* *		47	* *		39	* *	59	* *
Cambridge	27	* * * *	* *		20	* *		19	* *	27	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators ^(2, 3)										
Clinic	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)		Number	%	Number	%	Number	%
	Number	% (Rank)	% (Rank)	(Rank)						
Carroll Co	79	* * * *	* *	65	* *	58	* *	79	* *	
Cass Co	351	5.4 (11)	6.0 (41)	305	9.2 (8)	273	1.5 (14)	355	63.9 (71)	
Citizens	549	8.2 (60)	6.6 (51)	249	15.3 (75)	209	2.4 (34)	552	58.7 (87)	
Civic Ctr	896	9.2 (70)	6.1 (45)	773	15.3 (75)	651	2.6 (41)	902	66.1 (59)	
Clark Co	674	7.6 (47)	5.8 (35)	547	13.9 (62)	469	3.2 (61)	677	62.5 (73)	
Clay City	5	* * * *	* *	5	* *	5	* *	5	* *	
Clinton	87	* * * *	* *	71	* *	61	* *	87	* *	
Clinton Co	257	5.4 (11)	8.9 (93)	216	10.6 (19)	192	2.1 (28)	258	72.1 (38)	
Comm East	115	8.7 (65)	3.5 (5)	75	* *	60	* *	116	41.4 (101)	
Cottage Gr	793	9.1 (67)	7.2 (69)	585	13.8 (60)	501	3.2 (61)	801	56.2 (91)	
Crawford	76	* * * *	* *	61	* *	58	* *	76	* *	
Crown Pt	524	5.5 (13)	7.4 (76)	379	9.8 (13)	342	3.5 (67)	525	86.3 (2)	
Daviess Co	177	6.8 (31)	2.8 (3)	149	13.4 (59)	129	3.9 (72)	177	59.3 (86)	
Dearborn	308	9.4 (74)	9.4 (95)	241	12.9 (48)	208	3.4 (65)	312	64.7 (66)	

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators ^(2, 3)									
Clinic	Birthweight			Preterm ⁽⁶⁾		Full Term LBW ⁽⁷⁾		Ever Breastfed	
	Low ⁽⁴⁾	High ⁽⁵⁾		Number	%	Number	%	Number	%
	Number	% (Rank)	% (Rank)						
Decatur	155	6.5 (22)	5.2 (23)	133	12.8 (44)	116	4.3 (77)	155	82.6 (6)
Decatur Co	186	11.8 (96)	5.9 (39)	149	17.4 (87)	123	4.1 (74)	186	72.6 (33)
Dekalb Co	280	7.1 (35)	7.1 (65)	231	12.1 (36)	202	5.0 (88)	281	86.1 (3)
Dubois Co	242	4.5 (6)	5.8 (35)	187	12.8 (44)	163	0.6 (4)	242	76.9 (16)
E Chicago	535	10.7 (92)	5.8 (35)	428	16.8 (84)	355	2.5 (37)	537	64.4 (67)
Eaton	11	* * *	* *	8	* *	7	* *	11	* *
Elkhart	1,208	7.9 (52)	7.5 (78)	1,067	10.7 (21)	949	3.0 (52)	1,214	72.5 (35)
Elwood	85	* * *	* *	70	* *	58	* *	85	* *
Fayette Co	179	7.3 (42)	8.4 (91)	137	13.1 (51)	119	0.8 (8)	179	57.5 (89)
Floyd Co	447	9.4 (74)	5.8 (35)	367	19.3 (93)	293	2.4 (34)	450	65.6 (61)
Forest Man	664	11.4 (94)	5.1 (21)	498	17.9 (88)	407	2.7 (44)	670	60.9 (79)
Fountain	100	12.0 (98)	6.0 (41)	84	* *	69	* *	100	69.0 (52)
Franklin C	94	* * *	* *	78	* *	72	* *	94	* *
Fulton Co	129	3.1 (2)	7.8 (86)	107	9.3 (9)	97	* *	129	72.9 (30)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators ^(2, 3)									
Clinic	Birthweight			Preterm ⁽⁶⁾		Full Term LBW ⁽⁷⁾		Ever Breastfed	
	Low ⁽⁴⁾	High ⁽⁵⁾							
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Georgetown	1,314	9.7 (80)	6.9 (61)	932	15.1 (73)	789	3.0 (52)	1,317	72.6 (33)
Gibson Co	182	7.1 (35)	4.9 (17)	141	16.3 (83)	117	1.7 (17)	185	52.4 (94)
Glenn Pk	468	11.1 (93)	4.9 (17)	235	17.9 (88)	192	3.6 (70)	469	74.8 (23)
Goshen	576	6.3 (17)	10.1 (98)	489	12.1 (36)	428	2.1 (28)	581	79.7 (9)
Grant Co	527	10.2 (88)	5.5 (29)	425	13.2 (52)	367	4.9 (87)	529	50.1 (97)
Grant St	394	12.9(100)	4.8 (15)	304	18.8 (91)	245	4.5 (84)	396	51.3 (96)
Grassy Crk	915	10.1 (85)	4.8 (15)	678	15.9 (80)	568	2.8 (47)	918	66.4 (58)
Greene Co	159	8.8 (66)	6.3 (46)	127	11.0 (24)	113	4.4 (83)	159	64.2 (68)
Hamilton	649	8.5 (62)	7.1 (65)	509	12.2 (39)	446	3.1 (55)	650	77.5 (12)
Hammond	617	9.4 (74)	5.7 (32)	474	15.6 (78)	400	2.5 (37)	617	72.9 (30)
Hancock Co	283	6.4 (19)	7.4 (76)	218	11.9 (32)	192	0.5 (2)	283	76.7 (17)
Hanna	1,301	7.2 (40)	7.1 (65)	985	13.3 (55)	847	1.8 (18)	1,316	71.7 (40)
Harrison	235	6.4 (19)	6.8 (60)	187	16.0 (82)	157	1.9 (21)	235	74.0 (24)
Hendricks	364	6.6 (23)	6.6 (51)	264	9.8 (13)	238	1.3 (13)	364	76.4 (19)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators ^(2, 3)									
Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)						
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Henry Co	288	6.9 (33)	7.6 (81)	248	6.5 (2)	230	4.3 (77)	290	70.0 (48)
Hessville	351	6.6 (23)	7.7 (84)	279	10.8 (23)	248	3.2 (61)	353	66.9 (57)
Hispanic C	123	4.1 (3)	6.5 (48)	100	11.0 (24)	89	* *	123	84.6 (4)
Howard Co	657	6.7 (28)	5.2 (23)	546	10.1 (16)	489	2.7 (44)	660	51.5 (95)
Huntington	217	7.8 (51)	5.1 (21)	186	14.5 (68)	158	2.5 (37)	218	67.4 (55)
Jackson Co	326	7.4 (45)	8.3 (90)	274	15.0 (71)	230	3.0 (52)	328	72.0 (39)
Jasonville	7	* *	* *	6	* *	6	* *	7	* *
Jasper Co	206	4.4 (5)	10.2 (99)	160	8.8 (6)	146	0.7 (6)	206	60.7 (81)
Jay Co	148	10.1 (85)	2.0 (1)	110	11.8 (31)	97	* *	148	64.2 (68)
Jefferson	201	11.4 (94)	7.5 (78)	165	18.8 (91)	130	3.1 (55)	205	72.2 (37)
Jennings	260	8.1 (56)	6.5 (48)	223	14.8 (69)	190	0.5 (2)	260	67.3 (56)
Johnson Co	693	7.6 (47)	7.6 (81)	509	13.2 (52)	440	2.3 (31)	695	71.1 (41)
Kosciusko	410	6.6 (23)	6.6 (51)	317	12.3 (40)	278	0.7 (6)	411	73.5 (28)
Lagrange	130	9.2 (70)	6.9 (61)	104	15.4 (77)	88	* *	130	78.5 (10)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators ^(2, 3)									
Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)	High (5)		Number	%	Number	%	Number	%
	Number	% (Rank)	% (Rank)						
Lake Station	421	6.7 (28)	6.7 (57)	235	11.5 (30)	205	2.9 (49)	425	70.1 (47)
Laporte	386	4.9 (8)	6.5 (48)	318	10.7 (21)	284	1.8 (18)	386	73.3 (29)
Main St	791	9.9 (81)	4.4 (8)	633	13.0 (50)	551	4.5 (84)	791	60.4 (82)
Marshall	354	4.2 (4)	7.6 (81)	281	8.5 (5)	257	1.9 (21)	354	77.4 (13)
Martin Co	55	* * *	* * *	44	* *	34	* *	55	* *
Memorial	181	9.9 (81)	7.2 (69)	149	9.4 (10)	134	4.5 (84)	182	75.3 (22)
Merrillville	2	* * *	* * *	0	* *	0	* *	2	* *
Methodist	221	9.5 (77)	7.2 (69)	135	14.1 (66)	116	4.3 (77)	222	45.0(100)
Miami Co	185	8.6 (64)	5.9 (39)	160	15.0 (71)	136	1.5 (14)	185	59.5 (85)
Mich City	420	12.1 (99)	5.0 (20)	326	18.4 (90)	266	5.3 (89)	421	57.2 (90)
Moms/Kids	114	7.9 (52)	4.4 (8)	90	* *	78	* *	114	71.1 (41)
Monroe Co	589	8.0 (54)	9.8 (97)	468	11.3 (29)	414	3.1 (55)	590	80.8 (8)
Montgomery	258	5.8 (15)	11.6 (101)	207	7.2 (3)	191	2.6 (41)	259	69.9 (49)
Morgan Co	399	9.3 (73)	8.5 (92)	284	12.0 (35)	250	2.4 (34)	401	73.6 (27)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators ^(2, 3)									
Clinic	Birthweight			Preterm ⁽⁶⁾		Full Term LBW ⁽⁷⁾		Ever Breastfed	
	Low ⁽⁴⁾	High ⁽⁵⁾		Number	%	Number	%	Number	%
	Number	% ^(Rank)	% ^(Rank)						
N Arlingto	792	9.5 (77)	5.2 (23)	579	15.9 (80)	485	3.5 (67)	796	61.9 (74)
N Liberty	18	* *	* *	16	* *	16	* *	18	* *
N Manchest	47	* *	* *	37	* *	29	* *	47	* *
Newton Co	57	* *	* *	40	* *	32	* *	57	* *
Northwest	590	6.6 (23)	8.0 (87)	401	9.7 (12)	361	1.9 (21)	591	76.6 (18)
Ohio Co	6	* *	* *	4	* *	2	* *	7	* *
Orange Co	150	5.3 (9)	2.0 (1)	115	10.4 (17)	103	1.9 (21)	150	74.0 (24)
Owen Co	111	6.3 (17)	7.2 (69)	94	* *	79	* *	111	75.7 (21)
Parke Co	75	* *	* *	59	* *	51	* *	75	* *
Perry Co	122	13.9(101)	5.7 (32)	95	* *	83	* *	122	53.3 (93)
Pike Co	66	* *	* *	57	* *	41	* *	68	* *
Portage	368	7.6 (47)	7.3 (73)	293	11.9 (32)	257	2.3 (31)	369	70.2 (46)
Posey Co	93	* *	* *	71	* *	63	* *	93	* *
Pulaski Co	65	* *	* *	51	* *	44	* *	66	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators ^(2, 3)										
Clinic	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)							
	Number	% (Rank)	%	(Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Purdue	122	2.5 (1)	5.7 (32)		104	7.7 (4)	96	* *	122	94.3 (1)
Putnam Co	210	7.1 (35)	6.7 (57)		180	10.0 (15)	161	3.1 (55)	211	64.9 (64)
Randolph	166	6.0 (16)	6.6 (51)		143	13.3 (55)	124	0.0 (1)	167	61.1 (78)
Richmond	519	7.1 (35)	8.1 (88)		427	11.9 (32)	375	2.1 (28)	520	61.7 (77)
Ripley Co	176	9.1 (67)	6.3 (46)		136	13.2 (52)	118	0.8 (8)	176	61.9 (74)
Rush Co	97	* *	* *		84	* *	72	* *	99	* *
SB Ireland Rd	663	6.8 (31)	9.5 (96)		536	8.8 (6)	487	2.7 (44)	666	76.4 (19)
SB Lafayette	921	10.0 (84)	6.7 (57)		753	12.9 (48)	655	3.8 (71)	924	78.0 (11)
Scott Co	216	8.3 (61)	6.0 (41)		185	16.8 (84)	152	3.3 (64)	218	46.3 (99)
Shelby Co	304	7.2 (40)	4.9 (17)		241	13.3 (55)	207	4.3 (77)	306	67.6 (54)
Shelters	3	* *	* *		3	* *	1	* *	3	* *
SJ Reg Ctr	47	* *	* *		36	* *	34	* *	47	* *
So Calhoun	2,245	9.1 (67)	4.5 (11)		1,853	15.2 (74)	1,567	3.5 (67)	2,250	64.1 (70)
Spencer Co	96	* *	* *		79	* *	67	* *	96	* *

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
- (4) <2500 g.
- (5) >4000 g.
- (6) <37 weeks gestation.
- (7) 37 or more weeks gestation and <2500 g.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators ^(2, 3)										
Clinic	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)							
	Number	% (Rank)	% (Rank)		Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
St Francis	1	* * * *	* *		0	* *	0	* *	1	* *
St V-FLC	137	6.6 (23)	2.9 (4)		94	* *	84	* *	138	60.1 (83)
Starke Co	161	8.1 (56)	7.5 (78)		141	12.8 (44)	123	4.1 (74)	161	70.8 (44)
Steuben Co	253	7.5 (46)	11.5 (100)		224	11.2 (28)	196	3.1 (55)	256	83.6 (5)
Sullivan	149	8.1 (56)	5.4 (28)		132	10.6 (19)	118	4.2 (76)	149	69.8 (50)
Switz""Land	36	* * * *	* *		27	* *	26	* *	37	* *
Tiptecanoe	1,048	7.6 (47)	7.7 (84)		862	11.0 (24)	764	3.4 (65)	1,052	72.7 (32)
Tipton Co	55	* * * *	* *		40	* *	36	* *	55	* *
Union Co	45	* * * *	* *		33	* *	30	* *	45	* *
United Fai	325	9.2 (70)	5.5 (29)		264	14.0 (64)	227	2.6 (41)	325	69.5 (51)
Valpo	355	7.0 (34)	9.0 (94)		259	10.4 (17)	231	0.9 (11)	357	77.0 (14)
Vigo	575	8.0 (54)	7.3 (73)		475	12.8 (44)	407	1.2 (12)	583	65.4 (62)
Vincennes	179	7.3 (42)	4.5 (11)		152	13.8 (60)	131	3.1 (55)	180	62.8 (72)
W 21st Ave	320	10.3 (90)	4.7 (13)		256	16.8 (84)	211	4.3 (77)	322	58.1 (88)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators ^(2, 3)										
Clinic	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)							
	Number	% (Rank)	% (Rank)		Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Wabash	154	4.5 (6)	7.1 (65)		134	5.2 (1)	126	0.8 (8)	155	69.0 (52)
Warren Co	65	* * *	* *		48	* *	44	* *	65	* *
Warrick Co	228	9.6 (79)	7.0 (64)		191	15.7 (79)	160	0.6 (4)	229	65.1 (63)
Washington	148	8.1 (56)	8.1 (88)		128	12.5 (42)	112	1.8 (18)	148	72.3 (36)
Wells Co	148	10.1 (85)	4.1 (6)		116	12.1 (36)	102	2.9 (49)	148	77.0 (14)
Westside	646	7.3 (42)	6.0 (41)		397	14.4 (67)	339	1.5 (14)	650	60.8 (80)
White Co	193	6.7 (28)	7.3 (73)		163	11.0 (24)	143	2.8 (47)	196	65.8 (60)
Whitley Co	169	11.8 (96)	5.3 (26)		139	23.0 (94)	107	1.9 (21)	169	81.1 (7)
Wishard	123	10.6 (91)	4.1 (6)		84	* *	67	* *	123	61.8 (76)
Women""s	1	* * *	* *		1	* *	1	* *	1	* *
Unknown Clinics	2	* * *	* *		0	* *	0	* *	2	* *
Indiana	42,474	8.2	6.5		33,235	13.2	28,717	2.8	42,656	67.6
Nation (Prior Year)	880,933	8.0	6.7		948,286	10.7	749,107	3.4	871,811	68.2

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3A
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Racial and Ethnic Distribution by Clinic
Combined 3 Years

Comparison of Racial and Ethnic Distribution ⁽²⁾								
Clinic	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaska Native	Asian/Pacific Islander	Multiple Races	All Other/Unknown
		%	%	%	%	%	%	%
Action Ctr	120	26.7	58.3	10.0	0.0	0.0	4.2	0.8
Albion	826	76.9	0.4	20.6	0.1	0.8	0.6	0.6
Anderson	2,608	75.0	13.4	8.7	0.2	0.3	1.6	0.7
Ball Mem	22	*	*	*	*	*	*	*
Bartholome	1,632	80.3	1.8	15.3	0.0	0.6	1.1	0.9
Bedford	931	96.1	0.4	1.4	0.0	1.2	0.3	0.5
Benton Co	122	84.4	0.0	13.9	0.0	0.0	0.0	1.6
Berne	113	89.4	0.9	8.8	0.0	0.0	0.0	0.9
Bicknell	91	*	*	*	*	*	*	*
Blackburn	1,361	14.3	74.7	7.9	0.1	0.9	2.0	0.1
Blackford	293	96.9	0.3	1.7	0.0	0.0	0.3	0.7
Boone Co	612	94.0	1.1	2.9	0.0	0.7	0.2	1.1
Brazil	641	96.3	0.6	0.8	0.0	0.5	1.1	0.8
Brown Co	209	94.7	0.0	2.4	0.5	1.0	0.0	1.4
Cambridge	105	97.1	0.0	1.0	1.0	0.0	1.0	0.0
Carroll Co	248	89.1	0.0	10.1	0.0	0.4	0.4	0.0
Cass Co	1,148	63.6	0.7	32.7	0.0	1.0	0.9	1.1
Citizens	1,852	38.4	45.9	12.5	0.1	1.0	1.5	0.7
Civic Ctr	2,755	69.8	20.0	5.4	0.1	1.0	2.1	1.5
Clark Co	2,271	71.1	10.1	14.4	0.2	1.1	2.0	1.1

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3A
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Racial and Ethnic Distribution by Clinic
Combined 3 Years

Comparison of Racial and Ethnic Distribution ⁽²⁾								
Clinic	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaska Native	Asian/Pacific Islander	Multiple Races	All Other/Unknown
		%	%	%	%	%	%	%
Clay City	38	*	*	*	*	*	*	*
Clinton	303	98.0	0.3	0.0	0.0	0.3	0.3	1.0
Clinton Co	901	63.6	0.4	34.6	0.3	0.1	0.4	0.4
Comm East	330	31.8	60.6	4.2	0.0	0.6	2.7	0.0
Cottage Gr	2,538	72.1	9.9	15.1	0.1	0.5	1.7	0.7
Crawford	219	96.8	0.0	1.8	0.0	0.0	0.0	1.4
Crown Pt	1,729	81.1	6.4	9.5	0.1	0.9	1.2	0.8
Daviess Co	555	77.1	0.5	18.0	0.0	0.0	2.2	2.2
Dearborn	957	96.4	0.9	0.3	0.2	0.1	0.6	1.4
Decatur	506	93.1	0.4	5.5	0.0	0.0	0.2	0.8
Decatur Co	556	96.6	0.4	1.6	0.2	0.2	0.2	0.9
Dekalb Co	986	93.9	0.7	3.4	0.2	0.4	0.5	0.8
Dubois Co	782	73.0	0.0	23.7	0.1	1.7	0.6	0.9
E Chicago	1,852	6.3	36.2	56.4	0.1	0.2	0.3	0.5
Eaton	44	*	*	*	*	*	*	*
Elkhart	4,072	53.8	14.2	28.7	0.0	1.0	1.4	0.8
Elwood	312	93.3	0.6	4.5	0.0	0.6	0.0	1.0
Fayette Co	649	96.8	1.8	0.2	0.0	0.2	0.6	0.5
Floyd Co	1,410	76.0	12.5	6.1	0.1	2.2	1.3	1.8
Forest Man	2,229	11.5	81.1	4.3	0.2	0.9	1.0	1.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3A
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Racial and Ethnic Distribution by Clinic
Combined 3 Years

Comparison of Racial and Ethnic Distribution ⁽²⁾								
Clinic	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaska Native	Asian/Pacific Islander	Multiple Races	All Other/Unknown
		%	%	%	%	%	%	%
Fountain	356	95.5	0.0	2.2	0.0	0.8	0.3	1.1
Franklin C	325	98.5	0.0	0.3	0.0	0.0	0.3	0.9
Fulton Co	517	89.0	0.8	9.1	0.0	0.0	0.2	1.0
Georgetown	4,277	20.3	43.8	31.9	0.2	1.1	1.2	1.5
Gibson Co	578	91.5	3.3	2.4	0.7	0.7	1.0	0.3
Glenn Pk	1,894	17.1	68.4	10.6	0.2	0.8	2.1	0.8
Goshen	2,036	56.5	1.4	40.0	0.0	0.6	0.7	0.7
Grant Co	1,756	79.6	10.6	6.5	0.0	0.3	2.1	0.9
Grant St	1,207	2.3	90.6	4.8	0.2	0.4	0.9	0.7
Grassy Crk	3,217	25.1	53.1	18.4	0.2	0.8	1.9	0.5
Greene Co	522	97.1	0.4	1.7	0.0	0.4	0.2	0.2
Hamilton	2,047	77.8	6.9	11.0	0.2	2.5	1.3	0.3
Hammond	2,388	29.1	30.7	37.0	0.1	0.5	1.3	1.3
Hancock Co	866	94.3	1.4	1.8	0.1	0.2	0.8	1.3
Hanna	3,993	61.6	6.6	13.9	0.2	15.4	1.4	1.0
Harrison	727	93.3	0.3	3.9	0.0	0.4	1.0	1.2
Hendricks	1,273	86.6	4.6	4.1	0.1	3.0	0.1	1.5
Henry Co	1,003	94.5	0.8	2.1	0.1	0.3	1.2	1.0
Hessville	1,141	53.8	19.3	25.7	0.0	0.3	0.5	0.4
Hispanic C	370	3.8	9.7	83.2	0.3	0.3	0.0	2.7

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3A
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Racial and Ethnic Distribution by Clinic
Combined 3 Years

Comparison of Racial and Ethnic Distribution ⁽²⁾								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Howard Co	2,164	83.9	9.6	3.9	0.0	0.4	1.6	0.7
Huntington	726	96.3	0.6	1.5	0.0	0.3	0.7	0.7
IN0108 Closed	60	*	*	*	*	*	*	*
IN0904 Closed	326	90.2	4.0	1.5	0.0	0.6	1.5	2.1
IN2224 Closed	1	*	*	*	*	*	*	*
IN2305 Closed	14	*	*	*	*	*	*	*
Jackson Co	1,070	82.0	1.1	15.6	0.0	0.6	0.0	0.7
Jasonville	24	*	*	*	*	*	*	*
Jasper Co	697	87.8	0.4	10.9	0.0	0.1	0.4	0.3
Jay Co	466	89.5	0.6	7.9	0.0	0.2	0.0	1.7
Jefferson	733	87.2	1.8	7.8	0.0	0.5	1.5	1.2
Jennings	864	95.0	0.7	2.5	0.6	0.1	0.3	0.7
Johnson Co	2,227	87.4	1.4	5.4	0.0	3.1	1.1	1.5
Kosciusko	1,278	75.4	0.8	21.8	0.1	0.4	0.5	1.2
Lagrange	435	82.1	1.1	12.0	0.0	1.1	1.4	2.3
Lake Station	1,586	53.5	17.7	25.3	0.0	0.5	2.5	0.6
Laporte	1,381	80.1	3.3	14.8	0.0	0.3	0.9	0.7
Main St	2,555	81.7	12.3	2.5	0.2	0.7	2.2	0.5
Marshall	1,064	74.5	0.5	22.7	0.1	0.5	1.0	0.8
Martin Co	173	97.7	0.0	1.2	0.0	0.0	0.0	1.2

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3A
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Racial and Ethnic Distribution by Clinic
Combined 3 Years

Comparison of Racial and Ethnic Distribution ⁽²⁾								
Clinic	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaska Native	Asian/Pacific Islander	Multiple Races	All Other/Unknown
		%	%	%	%	%	%	%
Memorial	534	36.1	31.8	26.6	0.2	2.1	3.0	0.2
Merrillville	8	*	*	*	*	*	*	*
Methodist	812	31.7	51.4	14.0	0.1	2.2	0.6	0.0
Miami Co	670	91.8	1.8	2.2	0.6	0.7	2.8	0.0
Mich City	1,354	56.1	34.2	4.9	0.1	1.2	3.0	0.7
Moms/Kids	253	27.7	31.6	37.5	0.0	1.6	1.2	0.4
Monroe Co	2,024	81.1	5.7	3.9	0.0	5.9	2.1	1.3
Montgomery	872	87.6	0.8	9.4	0.1	0.3	0.9	0.8
Morgan Co	1,275	96.9	0.4	1.2	0.0	0.2	0.6	0.8
N Arlingto	2,448	35.6	44.9	15.4	0.0	0.9	2.4	0.7
N Liberty	51	*	*	*	*	*	*	*
N Manchest	181	87.8	0.6	6.1	0.0	0.0	5.0	0.6
Newton Co	185	92.4	0.0	7.0	0.0	0.5	0.0	0.0
Northwest	1,959	18.4	39.7	36.9	0.4	2.3	1.7	0.6
Ohio Co	29	*	*	*	*	*	*	*
Orange Co	487	94.7	1.6	1.6	0.0	0.8	0.8	0.4
Owen Co	388	95.6	0.3	1.8	0.0	1.0	0.5	0.8
Parke Co	240	96.7	0.0	1.7	0.4	0.0	0.4	0.8
Perry Co	386	97.2	0.5	0.8	0.0	0.3	0.0	1.3
Pike Co	226	96.0	0.4	1.8	0.0	0.0	1.3	0.4

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3A
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Racial and Ethnic Distribution by Clinic
Combined 3 Years

Comparison of Racial and Ethnic Distribution ⁽²⁾								
Clinic	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaska Native	Asian/Pacific Islander	Multiple Races	All Other/Unknown
		%	%	%	%	%	%	%
Portage	1,194	71.6	10.3	15.5	0.1	0.7	0.3	1.5
Posey Co	293	93.5	2.4	1.7	0.0	0.3	1.4	0.7
Pulaski Co	264	95.8	0.0	1.9	0.4	0.8	0.4	0.8
Purdue	427	32.3	4.2	6.8	1.2	52.0	1.4	2.1
Putnam Co	732	95.5	0.8	1.5	0.0	0.5	0.8	0.8
Randolph	622	88.3	0.2	7.4	0.2	0.5	1.4	2.1
Richmond	1,734	81.8	6.0	7.2	0.2	0.7	3.2	0.9
Ripley Co	610	93.3	0.0	3.6	0.7	0.0	1.0	1.5
Rush Co	312	95.8	0.3	1.6	0.6	0.6	0.6	0.3
SB Ireland Rd	2,385	71.6	16.6	7.2	0.3	1.3	2.2	0.9
SB Lafayette	2,786	32.7	38.4	22.6	0.3	1.9	2.6	1.5
Scott Co	720	96.0	0.0	2.2	0.0	0.4	0.7	0.7
Shelby Co	929	86.5	1.1	11.4	0.0	0.1	0.3	0.5
Shelters	24	*	*	*	*	*	*	*
SJ Reg Ctr	178	53.4	24.2	19.7	0.6	2.2	0.0	0.0
So Calhoun	7,405	46.6	28.7	15.4	0.1	6.4	2.2	0.6
Spencer Co	285	90.2	1.1	6.7	0.0	0.0	0.0	2.1
St Francis	10	*	*	*	*	*	*	*
St V-FLC	517	20.3	59.4	17.6	0.0	1.0	1.7	0.0
Starke Co	566	94.5	0.2	4.6	0.0	0.2	0.0	0.5

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3A
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Racial and Ethnic Distribution by Clinic
Combined 3 Years

Comparison of Racial and Ethnic Distribution ⁽²⁾								
Clinic	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaska Native	Asian/Pacific Islander	Multiple Races	All Other/Unknown
		%	%	%	%	%	%	%
Steuben Co	840	91.0	0.6	4.9	0.0	1.3	0.4	1.9
Sullivan	442	97.1	0.5	1.4	0.0	0.0	0.2	0.9
Switz""Land	148	94.6	1.4	0.7	0.0	0.0	0.7	2.7
Tiptecanoe	3,521	65.1	10.9	20.6	0.3	0.9	1.2	1.0
Tipton Co	182	91.8	0.5	7.1	0.0	0.0	0.5	0.0
Union Co	143	96.5	0.7	0.7	0.0	0.7	0.0	1.4
United Fai	1,241	76.2	14.4	5.4	0.0	0.8	2.2	1.0
Valpo	1,132	83.3	4.4	8.7	0.3	1.5	0.3	1.6
Vigo	2,070	86.6	8.7	2.0	0.0	0.9	1.4	0.5
Vincennes	649	93.4	0.6	1.7	0.2	0.3	1.5	2.3
W 21st Ave	1,229	2.0	95.4	1.5	0.1	0.0	0.5	0.6
Wabash	453	90.5	0.7	2.6	0.0	0.9	3.8	1.5
Warren Co	185	97.8	0.0	1.6	0.0	0.0	0.5	0.0
Warrick Co	777	95.2	1.4	1.0	0.0	0.3	1.3	0.8
Washington	544	96.1	0.2	1.8	0.4	0.4	0.7	0.4
Wells Co	461	92.4	0.4	4.3	0.2	0.2	1.3	1.1
Westside	2,316	41.8	21.2	33.8	0.2	0.9	1.1	0.9
White Co	668	78.7	0.0	19.6	0.1	0.4	0.4	0.6
Whitley Co	605	97.4	0.3	1.2	0.0	0.3	0.5	0.3
Wishard	655	13.4	49.0	35.0	0.0	1.4	1.1	0.2

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3A
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Racial and Ethnic Distribution by Clinic
Combined 3 Years

Comparison of Racial and Ethnic Distribution ⁽²⁾								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Women's	2	*	*	*	*	*	*	*
Unknown Clinics	308	68.5	17.2	9.1	0.0	1.9	1.0	2.3
Indiana	142,211	63.8	18.2	13.9	0.1	1.8	1.3	0.9

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4A
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Age Distribution by Clinic
Combined 3 Years

Comparison of Age Distribution ⁽²⁾								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Action Ctr	120	0.0	8.3	19.2	67.5	5.0	0.0	0.0
Albion	826	0.2	4.8	14.9	61.9	17.2	0.8	0.1
Anderson	2,608	0.4	5.6	14.7	63.7	14.5	1.0	0.1
Ball Mem	22	*	*	*	*	*	*	*
Bartholome	1,632	0.1	8.1	16.4	59.3	15.0	1.1	0.0
Bedford	931	0.2	9.0	17.0	61.0	11.6	1.2	0.0
Benton Co	122	0.0	6.6	11.5	63.9	16.4	1.6	0.0
Berne	113	0.9	7.1	12.4	60.2	19.5	0.0	0.0
Bicknell	91	*	*	*	*	*	*	*
Blackburn	1,361	0.4	6.0	11.6	61.5	19.3	1.2	0.0
Blackford	293	0.0	5.1	18.1	64.5	11.9	0.3	0.0
Boone Co	612	0.2	4.2	14.4	65.7	14.2	1.3	0.0
Brazil	641	0.2	8.6	15.3	62.7	12.3	0.9	0.0
Brown Co	209	0.0	6.2	15.8	59.8	15.3	2.9	0.0
Cambridge	105	0.0	7.6	14.3	61.0	16.2	1.0	0.0
Carroll Co	248	0.4	4.0	11.3	61.3	22.6	0.4	0.0
Cass Co	1,148	0.9	5.1	13.4	60.5	18.9	1.0	0.2
Citizens	1,852	0.2	5.7	13.6	60.1	19.0	1.2	0.2
Civic Ctr	2,755	0.3	6.0	13.8	64.8	14.6	0.5	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4A
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Age Distribution by Clinic
Combined 3 Years

Comparison of Age Distribution ⁽²⁾								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Clark Co	2,271	0.2	5.5	15.7	61.5	16.2	0.7	0.1
Clay City	38	*	*	*	*	*	*	*
Clinton	303	0.0	5.0	18.5	64.4	11.6	0.7	0.0
Clinton Co	901	0.0	6.5	13.2	63.2	15.8	1.3	0.0
Comm East	330	0.3	3.0	13.0	69.7	13.6	0.3	0.0
Cottage Gr	2,538	0.5	5.6	16.5	61.1	15.4	0.9	0.0
Crawford	219	0.0	5.5	17.8	64.8	11.9	0.0	0.0
Crown Pt	1,729	0.0	4.0	10.1	60.8	23.6	1.4	0.1
Daviess Co	555	0.0	5.0	13.3	65.9	13.9	0.7	1.1
Dearborn	957	0.5	4.5	15.0	64.3	14.3	1.0	0.3
Decatur	506	0.4	5.9	13.6	67.2	11.7	1.0	0.2
Decatur Co	556	0.0	8.3	15.8	60.4	14.7	0.7	0.0
Dekalb Co	986	0.3	5.2	16.4	64.3	12.8	0.9	0.1
Dubois Co	782	0.3	4.2	11.6	62.4	19.8	1.5	0.1
E Chicago	1,852	0.8	6.8	12.6	55.9	22.2	1.6	0.1
Eaton	44	*	*	*	*	*	*	*
Elkhart	4,072	0.5	6.0	13.6	60.2	18.5	1.2	0.0
Elwood	312	0.3	8.0	17.9	62.5	10.6	0.3	0.3
Fayette Co	649	0.3	9.6	16.6	60.9	11.9	0.6	0.2

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4A
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Age Distribution by Clinic
Combined 3 Years

Comparison of Age Distribution ⁽²⁾								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Floyd Co	1,410	0.4	6.3	16.2	61.8	14.6	0.4	0.3
Forest Man	2,229	0.5	5.3	13.8	63.0	16.1	1.3	0.0
Fountain	356	0.0	5.3	19.7	59.3	15.7	0.0	0.0
Franklin C	325	0.0	6.8	14.5	64.0	13.5	1.2	0.0
Fulton Co	517	0.2	4.8	19.3	60.3	14.3	1.0	0.0
Georgetown	4,277	0.3	4.9	10.7	58.5	24.2	1.3	0.1
Gibson Co	578	0.0	7.1	14.5	65.1	12.5	0.7	0.2
Glenn Pk	1,894	0.2	5.5	13.6	57.3	21.9	1.5	0.1
Goshen	2,036	0.3	5.4	14.6	55.6	22.4	1.6	0.0
Grant Co	1,756	0.7	7.5	15.9	63.9	11.4	0.6	0.1
Grant St	1,207	0.5	6.6	14.7	58.9	18.3	0.7	0.2
Grassy Crk	3,217	0.3	5.9	14.1	60.6	17.6	1.3	0.2
Greene Co	522	0.2	5.4	18.0	64.2	11.9	0.4	0.0
Hamilton	2,047	0.1	3.1	12.1	59.0	23.6	1.5	0.6
Hammond	2,388	0.3	6.1	12.8	59.0	19.9	1.8	0.1
Hancock Co	866	0.1	5.4	15.5	62.1	16.1	0.8	0.0
Hanna	3,993	0.2	4.3	10.4	64.7	19.3	1.1	0.1
Harrison	727	0.0	5.0	14.0	66.2	14.3	0.3	0.3
Hendricks	1,273	0.1	5.2	14.5	59.8	19.1	1.3	0.1

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4A
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Age Distribution by Clinic
Combined 3 Years

Comparison of Age Distribution ⁽²⁾								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Henry Co	1,003	0.6	6.0	15.5	64.2	13.2	0.5	0.1
Hessville	1,141	0.4	5.3	12.4	58.6	21.7	1.5	0.2
Hispanic C	370	1.1	5.1	6.2	58.9	27.6	1.1	0.0
Howard Co	2,164	0.3	5.6	15.7	63.2	14.1	0.8	0.3
Huntington	726	0.1	3.9	19.1	64.7	10.3	0.4	1.4
IN0108 Closed	60	*	*	*	*	*	*	*
IN0904 Closed	326	0.9	8.6	15.0	62.9	12.0	0.6	0.0
IN2224 Closed	1	*	*	*	*	*	*	*
IN2305 Closed	14	*	*	*	*	*	*	*
Jackson Co	1,070	0.1	8.4	14.9	63.0	12.7	0.7	0.3
Jasonville	24	*	*	*	*	*	*	*
Jasper Co	697	0.0	5.7	15.2	60.1	17.1	1.7	0.1
Jay Co	466	0.0	6.2	14.4	64.2	14.8	0.4	0.0
Jefferson	733	0.5	6.7	16.1	61.8	14.1	0.8	0.0
Jennings	864	0.2	8.9	21.1	56.4	13.1	0.3	0.0
Johnson Co	2,227	0.1	5.9	14.8	61.9	16.1	1.2	0.0
Kosciusko	1,278	0.3	6.7	15.2	58.2	18.4	1.3	0.0
Lagrange	435	0.2	6.0	13.6	58.6	19.1	1.8	0.7
Lake Station	1,586	0.3	5.2	12.7	59.8	20.9	1.0	0.1

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4A
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Age Distribution by Clinic
Combined 3 Years

Comparison of Age Distribution ⁽²⁾								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Laporte	1,381	0.1	4.9	12.7	61.7	19.6	1.0	0.0
Main St	2,555	0.2	5.6	14.3	64.9	14.0	0.9	0.1
Marshall	1,064	0.1	5.3	12.1	57.7	22.7	1.9	0.3
Martin Co	173	0.0	9.8	13.9	61.8	13.9	0.6	0.0
Memorial	534	0.2	6.0	12.9	59.9	19.7	1.3	0.0
Merrillville	8	*	*	*	*	*	*	*
Methodist	812	0.2	5.4	15.1	63.8	14.7	0.7	0.0
Miami Co	670	0.4	6.0	17.8	60.7	14.5	0.4	0.1
Mich City	1,354	0.5	6.1	14.3	59.5	18.2	1.4	0.0
Moms/Kids	253	0.0	2.4	13.0	57.3	25.3	2.0	0.0
Monroe Co	2,024	0.1	3.3	13.7	63.2	17.8	1.4	0.6
Montgomery	872	0.3	6.7	15.3	60.2	16.1	1.5	0.0
Morgan Co	1,275	0.2	5.8	14.6	64.5	13.9	0.9	0.2
N Arlingto	2,448	0.3	6.3	11.8	61.8	18.8	1.0	0.0
N Liberty	51	*	*	*	*	*	*	*
N Manchest	181	0.0	4.4	16.0	66.9	12.2	0.6	0.0
Newton Co	185	0.5	3.2	13.0	65.9	16.8	0.5	0.0
Northwest	1,959	0.1	3.4	8.8	63.3	22.8	1.6	0.0
Ohio Co	29	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4A
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Age Distribution by Clinic
Combined 3 Years

Comparison of Age Distribution ⁽²⁾								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Orange Co	487	0.6	4.7	15.6	61.6	17.0	0.4	0.0
Owen Co	388	0.3	5.4	17.3	63.4	12.1	1.5	0.0
Parke Co	240	0.0	8.8	18.8	58.3	13.3	0.8	0.0
Perry Co	386	0.5	8.0	16.6	61.4	13.0	0.5	0.0
Pike Co	226	0.0	6.2	17.3	61.9	14.2	0.4	0.0
Portage	1,194	0.0	4.2	11.6	63.1	19.5	1.3	0.3
Posey Co	293	0.0	6.5	16.7	62.1	14.0	0.7	0.0
Pulaski Co	264	0.4	4.2	17.0	64.4	12.9	0.8	0.4
Purdue	427	0.0	0.9	3.7	56.4	36.5	1.6	0.7
Putnam Co	732	0.5	6.6	15.7	65.0	11.5	0.7	0.0
Randolph	622	0.3	5.1	14.8	63.0	15.8	0.6	0.3
Richmond	1,734	0.2	5.4	14.6	64.4	14.8	0.6	0.0
Ripley Co	610	0.0	6.4	16.1	61.5	15.2	0.8	0.0
Rush Co	312	0.6	7.4	16.3	64.7	10.3	0.6	0.0
SB Ireland Rd	2,385	0.2	4.9	13.3	61.1	19.2	1.2	0.0
SB Lafayette	2,786	0.3	7.5	13.9	57.5	19.4	1.4	0.0
Scott Co	720	0.4	6.3	16.8	66.3	9.3	1.0	0.0
Shelby Co	929	0.2	6.4	15.5	63.0	13.8	1.2	0.0
Shelters	24	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4A
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Age Distribution by Clinic
Combined 3 Years

Comparison of Age Distribution ⁽²⁾								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
SJ Reg Ctr	178	0.0	4.5	15.2	57.3	22.5	0.6	0.0
So Calhoun	7,405	0.3	4.6	12.9	63.1	17.4	1.2	0.6
Spencer Co	285	0.4	5.6	14.7	61.8	16.5	1.1	0.0
St Francis	10	*	*	*	*	*	*	*
St V-FLC	517	0.2	2.3	10.3	67.7	18.4	1.2	0.0
Starke Co	566	0.9	5.3	14.3	61.5	16.6	1.2	0.2
Steuben Co	840	0.1	5.1	16.2	64.4	13.5	0.7	0.0
Sullivan	442	0.7	8.4	18.8	57.7	13.8	0.7	0.0
Switz""Land	148	0.0	7.4	14.2	60.8	14.9	2.7	0.0
Tippecanoe	3,521	0.2	6.2	14.3	60.0	18.0	1.2	0.1
Tipton Co	182	0.0	11.0	13.7	61.0	13.2	1.1	0.0
Union Co	143	0.0	5.6	16.8	58.0	18.2	1.4	0.0
United Fai	1,241	0.1	4.0	12.8	64.1	17.6	0.4	1.0
Valpo	1,132	0.1	4.9	12.4	61.9	19.1	1.6	0.1
Vigo	2,070	0.4	8.0	16.2	62.3	12.2	0.8	0.0
Vincennes	649	0.3	6.2	16.9	61.9	13.9	0.5	0.3
W 21st Ave	1,229	0.8	9.0	15.0	55.7	18.5	0.8	0.2
Wabash	453	0.2	4.6	14.6	67.3	11.3	1.5	0.4
Warren Co	185	0.0	6.5	18.9	60.0	14.1	0.5	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4A
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Age Distribution by Clinic
Combined 3 Years

Comparison of Age Distribution ⁽²⁾								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Warrick Co	777	0.0	4.5	15.2	64.7	15.3	0.0	0.3
Washington	544	0.0	7.2	18.6	62.1	10.8	0.9	0.4
Wells Co	461	0.7	4.1	14.5	65.5	14.5	0.7	0.0
Westside	2,316	0.4	5.9	11.3	60.2	20.9	1.3	0.0
White Co	668	0.7	6.9	13.5	60.6	16.3	1.9	0.0
Whitley Co	605	0.3	4.3	15.4	65.8	12.9	1.3	0.0
Wishard	655	0.6	5.8	13.0	55.4	22.6	2.6	0.0
Women""s	2	*	*	*	*	*	*	*
Unknown Clinics	308	0.3	6.5	11.7	65.6	14.6	1.3	0.0
Indiana	142,211	0.3	5.7	14.0	61.6	17.3	1.1	0.1

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Maternal Health Indicators by Clinic
Combined 3 Years

Comparison of Maternal Health Indicators ^(2, 3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight	Over- weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Action Ctr	88	* *	* *	81	* *	* *	23	* *	16	* *	113	4.4(100)	113	7.1(111)
Albion	650	4.5 (38)	52.6 (45)	611	17.8 (46)	53.0 (50)	146	39.0 (37)	161	31.1 (63)	788	0.5 (10)	788	1.6(16)
Anderson	1,933	4.3 (29)	55.6 (82)	1,850	16.5 (18)	53.9 (59)	478	45.4 (54)	523	32.9(70)	2,538	1.8 (35)	2,538	2.5(43)
Ball Mem	12	* *	* *	12	* *	* *	2	* *	0	* *	22	* *	22	* *
Bartholome	1,291	7.5 (116)	52.9 (49)	1,240	16.9 (26)	55.5 (84)	259	32.4 (20)	250	26.4(43)	1,590	5.3(116)	1,590	6.6(106)
Bedford	746	6.4 (99)	53.6 (61)	704	17.6 (39)	51.4 (28)	103	49.5 (59)	176	32.4(69)	901	4.9(111)	901	7.7(112)
Benton Co	96	* *	* *	92	* *	* *	15	* *	26	* *	118	7.6(124)	118	5.1(81)
Berne	94	* *	* *	92	* *	* *	19	* *	28	* *	112	1.8 (35)	112	1.8(21)
Bicknell	75	* *	* *	69	* *	* *	16	* *	48	* *	86	* *	86	* *
Blackburn	950	4.7 (47)	55.5 (79)	892	18.7 (65)	54.3 (67)	227	44.9 (53)	220	30.9(62)	1,313	4.7(107)	1,313	8.0(115)
Blackford	237	6.3 (97)	49.4 (13)	231	19.5 (89)	48.9 (11)	56	* *	46	* *	289	3.1 (66)	289	9.0(121)
Boone Co	464	6.7 (104)	51.7 (32)	438	17.4 (36)	51.6 (31)	85	* *	155	21.3(22)	586	1.0 (18)	586	1.2(15)
Brazil	536	5.4 (72)	56.9 (96)	511	19.8 (96)	54.0 (60)	75	* *	112	17.9(10)	623	2.9 (64)	623	3.7(60)
Brown Co	167	6.0 (88)	43.7 (2)	159	18.9 (71)	50.9 (22)	32	* *	56	* *	202	7.9(126)	202	5.9(95)
Cambridge	79	* *	* *	75	* *	* *	16	* *	18	* *	101	2.0 (40)	101	5.9(95)
Carroll Co	199	3.0 (6)	56.3 (91)	193	17.6 (39)	48.2 (8)	34	* *	43	* *	246	0.8 (15)	246	2.0(26)
Cass Co	917	4.6 (41)	55.0 (74)	845	20.6(103)	48.2 (8)	171	29.8 (14)	135	20.0(14)	1,082	3.3 (73)	1,082	1.9(22)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Maternal Health Indicators by Clinic
Combined 3 Years

Comparison of Maternal Health Indicators ^(2, 3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Citizens	1,108	4.3 (29)	57.2 (101)	887	18.7 (65)	55.4 (80)	246	47.2 (56)	338	34.9(74)	1,635	3.9 (87)	1,635	6.3(101)
Civic Ctr	2,293	4.2 (25)	56.1 (89)	1,945	18.3 (57)	57.2(106)	454	39.0 (37)	1,569	24.5(37)	2,421	2.7 (59)	2,421	4.1(68)
Clark Co	1,784	4.6 (41)	56.8 (95)	1,693	19.2 (82)	52.4 (44)	367	34.6 (27)	425	31.3(65)	2,188	6.5(123)	2,188	2.5(43)
Clay City	32	* *	* *	30	* *	* *	4	* *	6	* *	37	* *	37	* *
Clinton	249	7.2 (112)	49.0 (8)	240	16.7 (23)	56.3 (97)	61	* *	43	* *	297	4.4(100)	297	6.1(98)
Clinton Co	753	3.9 (19)	62.8 (121)	724	19.3 (83)	49.9 (15)	171	33.9 (23)	155	23.2(29)	882	2.3 (47)	882	2.6(49)
Comm East	202	6.4 (99)	53.5 (60)	198	17.2 (33)	59.1(119)	40	* *	0	* *	328	4.0 (92)	328	15.2(127)
Cottage Gr	1,855	4.9 (57)	55.5 (79)	1,714	19.0 (72)	54.5 (69)	404	30.7 (15)	488	23.2(29)	2,413	3.7 (81)	2,413	5.4(89)
Crawford	175	5.1 (61)	57.1 (97)	166	20.5(101)	51.8 (35)	38	* *	34	* *	212	1.9 (38)	212	2.4(41)
Crown Pt	1,254	4.8 (51)	49.1 (10)	1,183	14.7 (6)	56.6(102)	301	24.3 (4)	369	16.5 (6)	1,664	4.1 (95)	1,664	3.8(64)
Daviess Co	444	2.9 (5)	61.0 (117)	400	20.8(108)	50.5 (19)	92	* *	315	37.8(78)	514	2.5 (53)	514	8.0(115)
Dearborn	756	5.8 (83)	48.9 (7)	719	14.5 (5)	57.6(110)	146	48.6 (58)	210	23.3(32)	922	1.5 (28)	922	2.2(33)
Decatur	403	6.2 (96)	52.9 (49)	382	16.5 (18)	56.3 (97)	63	* *	101	27.7(53)	487	0.4 (7)	487	2.1(30)
Decatur Co	434	6.7 (104)	55.8 (83)	419	17.4 (36)	51.3 (27)	53	* *	124	14.5 (3)	540	3.9 (87)	540	5.9(95)
Dekalb Co	788	5.2 (63)	50.6 (19)	746	19.3 (83)	54.6 (72)	160	28.8 (8)	130	23.8(34)	953	0.9 (16)	953	2.3(40)
Dubois Co	589	5.1 (61)	61.1 (118)	570	17.7 (43)	53.5 (56)	135	26.7 (5)	133	17.3 (8)	767	5.0(112)	767	3.1(54)
E Chicago	1,439	2.8 (2)	62.8 (121)	1,378	20.7(106)	51.9 (36)	290	50.0 (61)	158	39.9(82)	1,805	4.2 (98)	1,805	6.9(109)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Maternal Health Indicators by Clinic
Combined 3 Years

Comparison of Maternal Health Indicators ^(2, 3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Eaton	31	* *	* *	30	* *	* *	6	* *	2	* *	42	* *	42	* *
Elkhart	3,434	4.0 (22)	57.4 (105)	3,267	21.0(110)	51.2 (26)	408	29.2 (11)	642	20.4 (16)	3,929	2.3 (47)	3,929	3.7(60)
Elwood	236	8.1 (119)	53.4 (58)	226	23.9(119)	46.9 (4)	51	* *	64	* *	305	2.6 (56)	305	2.6(49)
Fayette Co	525	7.2 (112)	52.0 (36)	504	17.3 (34)	58.1(115)	87	* *	105	33.3(72)	631	1.1 (20)	631	2.1(30)
Floyd Co	1,093	5.3 (67)	53.8 (62)	1,019	19.4 (86)	52.3 (41)	201	34.8 (28)	409	25.4(41)	1,342	8.3(127)	1,342	12.4(126)
Forest Man	1,626	4.1 (23)	60.1 (114)	1,487	18.6 (63)	56.5(100)	335	39.4 (40)	286	27.3(51)	2,104	2.6 (56)	2,104	4.1(68)
Fountain	287	2.4 (1)	55.1 (75)	273	19.0 (72)	60.4(121)	52	* *	63	* *	343	5.2(114)	343	5.5(91)
Franklin C	249	5.2 (63)	57.4 (105)	237	19.0 (72)	55.7 (90)	40	* *	59	* *	316	1.9 (38)	316	2.2(33)
Fulton Co	423	4.3 (29)	52.2 (41)	390	18.5 (61)	58.2(116)	71	* *	113	28.3(54)	491	0.0 (1)	491	0.2(3)
Georgetown	3,032	3.9 (19)	57.2 (101)	2,813	18.4 (58)	52.5 (45)	754	43.2 (50)	740	31.4(66)	4,077	4.5(102)	4,077	5.1(81)
Gibson Co	437	5.3 (67)	53.1 (54)	407	18.7 (65)	55.5 (84)	81	* *	101	22.8(26)	549	1.6 (29)	549	2.0(26)
Glenn Pk	1,129	4.3 (29)	57.8 (108)	811	19.0 (72)	54.0 (60)	270	48.5 (57)	197	24.9(39)	1,576	1.4 (27)	1,576	3.0(53)
Goshen	1,689	3.5 (11)	57.3 (104)	1,616	16.9 (26)	53.1 (51)	215	23.3 (3)	327	21.1(20)	1,968	4.5(102)	1,968	2.0(26)
Grant Co	1,419	5.9 (86)	51.4 (29)	1,337	19.3 (83)	53.3 (54)	279	45.9 (55)	487	26.9(48)	1,684	2.7 (59)	1,684	8.3(118)
Grant St	927	3.9 (19)	60.9 (116)	867	22.0(115)	55.4 (80)	203	57.1 (62)	114	50.0(83)	1,171	0.5 (10)	1,171	3.1(54)
Grassy Crk	2,161	4.2 (25)	59.3 (113)	1,914	17.7 (43)	55.4 (80)	441	36.3 (31)	463	28.7(55)	2,978	2.3 (47)	2,978	4.1(68)
Greene Co	411	6.1 (92)	52.1 (37)	390	20.8(108)	55.6 (87)	82	* *	121	24.0(36)	509	3.1 (66)	509	1.0(13)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Maternal Health Indicators by Clinic
Combined 3 Years

Comparison of Maternal Health Indicators ^(2, 3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight	Over- weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
							Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)
Hamilton	1,539	4.5 (38)	51.3 (27)	1,465	19.1 (79)	54.1 (63)	384	40.6 (45)	549	31.5(67)	1,984	3.9 (87)	1,984	4.1(68)
Hammond	1,648	4.1 (23)	58.7 (111)	1,456	18.1 (52)	54.1 (63)	359	44.3 (52)	332	37.3(76)	2,199	2.0 (40)	2,199	2.5(43)
Hancock Co	651	5.7 (79)	52.1 (37)	614	14.7 (6)	59.1(119)	149	28.9 (9)	193	16.1 (5)	834	3.5 (75)	834	2.4(41)
Hanna	2,843	5.0 (58)	50.7 (21)	2,638	22.4(116)	48.8 (10)	598	31.3 (17)	697	23.2(29)	3,814	4.3 (99)	3,814	4.2(72)
Harrison	565	5.0 (58)	52.9 (49)	528	21.8(114)	50.0 (16)	78	* *	101	17.8 (9)	694	2.7 (59)	694	2.6(49)
Hendricks	942	4.7 (47)	52.1 (37)	801	12.2 (2)	57.7(112)	248	41.5 (47)	674	39.6(81)	1,136	2.4 (51)	1,136	2.5(43)
Henry Co	828	6.8 (109)	50.7 (21)	797	18.2 (56)	57.5(109)	127	31.5 (19)	127	18.9(12)	979	1.8 (35)	979	0.9(12)
Hessville	871	3.8 (16)	57.2 (101)	833	17.9 (48)	57.9(113)	186	36.6 (32)	146	30.8(60)	1,108	0.7 (13)	1,108	1.0(13)
Hispanic C	287	2.8 (2)	57.1 (97)	268	29.1(123)	38.8 (2)	63	* *	48	* *	351	3.1 (66)	351	3.7(60)
Howard Co	1,761	5.2 (63)	51.8 (35)	1,699	19.4 (86)	50.9 (22)	309	41.7 (48)	373	32.2(68)	2,106	1.6 (29)	2,106	2.2(33)
Huntington	624	8.3 (120)	49.7 (15)	596	19.5 (89)	54.5 (69)	75	* *	91	* *	702	3.8 (86)	702	9.7(123)
IN0108 Closed	51	* *	* *	49	* *	* *	12	* *	13	* *	58	* *	58	* *
IN0904 Closed	283	6.7 (104)	49.1 (10)	229	19.7 (94)	52.8 (48)	59	* *	232	12.9 (2)	269	4.1 (95)	269	4.8(77)
IN2224 Closed	1	* *	* *	1	* *	* *	0	* *	0	* *	1	* *	1	* *
IN2305 Closed	13	* *	* *	8	* *	* *	2	* *	5	* *	9	* *	9	* *
Jackson Co	896	6.0 (88)	54.9 (73)	862	19.1 (79)	52.1 (38)	167	38.3 (35)	218	23.9(35)	1,047	3.2 (71)	1,047	5.3(87)
Jasonville	21	* *	* *	20	* *	* *	3	* *	5	* *	24	* *	24	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Maternal Health Indicators by Clinic
Combined 3 Years

Comparison of Maternal Health Indicators ^(2, 3)														
Clinic	Prepregnancy BMI ⁽⁴⁾			Weight Gain ⁽⁵⁾			Anemia (Low Hb/Hct) ⁽⁶⁾				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Jasper Co	540	4.8 (51)	50.0 (16)	492	16.5 (18)	53.7 (58)	60	* *	95	* *	652	0.3 (6)	652	0.5(7)
Jay Co	363	5.8 (83)	55.1 (75)	345	19.7 (94)	53.3 (54)	69	* *	81	* *	450	3.1 (66)	450	2.2(33)
Jefferson	581	5.7 (79)	54.2 (68)	547	16.3 (16)	52.3 (41)	100	31.0 (16)	126	11.1 (1)	704	5.4(117)	704	6.3(101)
Jennings	756	6.7 (104)	53.0 (53)	727	19.5 (89)	51.6 (31)	77	* *	118	28.8(56)	840	1.0 (18)	840	1.7(20)
Johnson Co	1,628	6.1 (92)	51.1 (26)	1,546	16.5 (18)	56.0 (92)	377	35.5 (30)	424	20.8(18)	2,156	3.5 (75)	2,156	7.9(114)
Kosciusko	1,001	4.7 (47)	55.5 (79)	926	17.8 (46)	53.1 (51)	245	33.9 (23)	336	21.4(23)	1,205	4.5(102)	1,205	5.0(79)
Lagrange	337	5.3 (67)	50.1 (17)	316	18.4 (58)	55.1 (75)	74	* *	122	23.0(27)	412	4.1 (95)	412	6.1(98)
Lake Station	1,001	5.8 (83)	54.0 (66)	809	16.6 (22)	58.0(114)	174	39.1 (39)	161	21.1(20)	1,399	3.9 (87)	1,399	6.2(100)
Laporte	1,132	5.7 (79)	52.2 (41)	1,047	16.2 (13)	54.3 (67)	238	20.6 (1)	466	18.7(11)	1,304	4.0 (92)	1,304	1.6(16)
Main St	1,933	5.5 (75)	54.0 (66)	1,818	19.0 (72)	55.9 (91)	380	40.3 (42)	309	31.1(63)	2,451	2.1 (44)	2,451	2.1(30)
Marshall	856	3.2 (8)	61.1(118)	814	16.3 (16)	56.3 (97)	187	23.0 (2)	165	17.0 (7)	1,029	5.7(120)	1,029	6.5(105)
Martin Co	143	4.2 (25)	51.0 (25)	137	11.7 (1)	62.0(122)	21	* *	14	* *	166	1.2 (22)	166	5.4(89)
Memorial	450	6.7 (104)	56.0 (87)	445	20.0 (98)	46.7 (3)	84	* *	1	* *	531	1.3 (26)	531	0.8(11)
Merrillville	3	* *	* *	0	* *	* *	0	* *	0	* *	5	* *	5	* *
Methodist	568	5.3 (67)	56.0 (87)	559	19.5 (89)	55.5 (84)	100	34.0 (25)	1	* *	806	5.2(114)	806	10.4(124)
Miami Co	556	6.5 (101)	52.3 (43)	520	17.7 (43)	55.2 (76)	95	* *	401	25.9(42)	637	2.8 (62)	637	5.2(84)
Mich City	1,014	4.6 (41)	57.1 (97)	915	16.0 (12)	55.3 (79)	206	49.5 (59)	327	37.3(76)	1,263	0.9 (16)	1,263	2.7(52)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Maternal Health Indicators by Clinic
Combined 3 Years

Comparison of Maternal Health Indicators ^(2, 3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight	Over- weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
							Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)
Moms/Kids	183	3.3 (10)	65.6 (124)	161	13.7 (4)	54.0 (60)	39	* *	31	* *	231	2.6 (56)	231	5.2(84)
Monroe Co	1,628	5.6 (77)	49.4 (13)	1,520	17.9 (48)	52.3 (41)	297	29.3 (13)	464	24.6(38)	1,919	2.4 (51)	1,919	1.9(22)
Montgomery	717	6.1 (92)	52.6 (45)	681	17.9 (48)	57.6(110)	140	28.6 (6)	194	21.6(25)	843	3.7 (81)	843	4.7(75)
Morgan Co	926	6.6 (103)	55.3 (77)	872	17.3 (34)	55.2 (76)	173	40.5 (44)	224	30.8(60)	1,228	1.7 (32)	1,228	2.2(33)
N Arlingto	1,804	4.6 (41)	58.9 (112)	1,682	17.0 (28)	56.5(100)	358	38.5 (36)	317	25.2(40)	2,335	3.0 (65)	2,335	5.6(93)
N Liberty	41	* *	* *	40	* *	* *	6	* *	12	* *	50	* *	50	* *
N Manchest	148	8.8 (122)	51.4 (29)	138	18.1 (52)	50.7 (21)	26	* *	117	15.4 (4)	172	4.7(107)	172	5.8(94)
Newton Co	141	5.7 (79)	48.2 (5)	136	17.6 (39)	47.1 (5)	23	* *	26	* *	180	0.0 (1)	180	0.6(8)
Northwest	1,332	3.5 (11)	57.1 (97)	1,242	18.0 (51)	52.2 (39)	328	36.9 (33)	320	27.5(52)	1,877	3.7 (81)	1,877	3.7(60)
Ohio Co	22	* *	* *	22	* *	* *	5	* *	2	* *	29	* *	29	* *
Orange Co	411	4.4 (35)	60.6 (115)	393	22.6(117)	48.1 (7)	52	* *	56	* *	471	4.5(102)	471	11.7(125)
Owen Co	334	7.2 (112)	52.1 (37)	323	17.0 (28)	51.7 (34)	51	* *	78	* *	377	6.4(122)	377	6.4(103)
Parke Co	199	6.5 (101)	49.2 (12)	197	16.2 (13)	56.9(105)	20	* *	37	* *	240	1.7 (32)	240	2.5(43)
Perry Co	289	5.2 (63)	46.7 (3)	276	19.6 (93)	50.0 (16)	59	* *	72	* *	374	3.5 (75)	374	5.3(87)
Pike Co	175	4.6 (41)	54.3 (69)	165	18.8 (69)	58.2(116)	31	* *	26	* *	216	3.7 (81)	216	8.3(118)
Portage	918	4.4 (35)	52.7 (47)	858	17.4 (36)	56.2 (95)	178	28.7 (7)	214	29.0(57)	1,141	3.3 (73)	1,141	3.9(65)
Posey Co	226	5.3 (67)	55.8 (83)	207	15.9 (11)	55.6 (87)	52	* *	57	* *	275	0.4 (7)	275	0.4(6)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Maternal Health Indicators by Clinic
Combined 3 Years

Comparison of Maternal Health Indicators ^(2, 3)														
Clinic	Prepregnancy BMI ⁽⁴⁾			Weight Gain ⁽⁵⁾			Anemia (Low Hb/Hct) ⁽⁶⁾				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Pulaski Co	199	3.0 (6)	50.8 (23)	178	18.5 (61)	55.6 (87)	24	* *	27	* *	243	0.4 (7)	243	0.0(1)
Purdue	353	7.9 (118)	25.8 (1)	338	21.6(113)	34.9 (1)	53	* *	76	* *	413	0.7 (13)	413	0.2(3)
Putnam Co	605	4.3 (29)	53.9 (65)	572	16.8 (25)	58.6(118)	80	* *	158	29.1(58)	706	5.8(121)	706	8.8(120)
Randolph	501	9.0(124)	53.1 (54)	478	18.6 (63)	54.2 (65)	115	37.4 (34)	101	26.7(44)	605	1.2 (22)	605	4.5(74)
Richmond	1,442	5.5 (75)	55.3 (77)	1,368	22.9(118)	50.2 (18)	208	31.3 (17)	278	35.3(75)	1,682	5.5(118)	1,682	3.9(65)
Ripley Co	467	4.3 (29)	56.5 (92)	437	18.1 (52)	54.7 (73)	72	* *	119	38.7(80)	582	5.0(112)	582	9.5(122)
Rush Co	253	7.1 (111)	49.0 (8)	244	18.4 (58)	54.5 (69)	37	* *	72	* *	305	3.9 (87)	305	5.2(84)
SB Ireland Rd	1,856	5.0 (58)	54.4 (72)	1,737	15.8 (10)	57.4(108)	356	34.3 (26)	592	30.4(59)	2,275	2.0 (40)	2,275	3.5(59)
SB Lafayette	2,211	3.8 (16)	56.7 (94)	2,080	20.5(101)	51.5 (30)	426	43.4 (51)	679	33.1(71)	2,676	1.2 (22)	2,676	1.6(16)
Scott Co	597	7.4 (115)	53.1 (54)	573	23.9(119)	51.1 (24)	75	* *	123	26.8(46)	697	3.6 (79)	697	4.7(75)
Shelby Co	703	6.0 (88)	54.3 (69)	665	17.1 (31)	56.1 (93)	134	40.3 (42)	423	27.0(49)	894	2.0 (40)	894	1.9(22)
Shelters	17	* *	* *	14	* *	* *	2	* *	3	* *	21	* *	21	* *
SJ Reg Ctr	146	4.8 (51)	53.4 (58)	146	17.1 (31)	56.2 (95)	34	* *	0	* *	178	0.0 (1)	178	0.0(1)
So Calhoun	6,066	4.5 (38)	54.3 (69)	5,791	20.7(106)	50.5 (19)	1,075	42.0 (49)	1,059	33.4(73)	7,178	2.1 (44)	7,178	3.4(58)
Spencer Co	223	7.6 (117)	53.8 (62)	215	18.1 (52)	49.3 (13)	53	* *	30	* *	281	2.5 (53)	281	3.2(56)
St Francis	0	* *	* *	0	* *	* *	0	* *	0	* *	10	* *	10	* *
St V-FLC	369	4.6 (41)	61.8 (120)	353	19.0 (72)	55.2 (76)	92	* *	12	* *	504	2.8 (62)	504	4.8(77)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Maternal Health Indicators by Clinic
Combined 3 Years

Comparison of Maternal Health Indicators ^(2, 3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight	Over- weight		< Ideal	> Ideal	3rd Trimester		Postpartum						
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Starke Co	466	5.4 (72)	50.9 (24)	446	15.7 (9)	56.1 (93)	60	* *	64	* *	550	2.5 (53)	550	4.4(73)
Steuben Co	704	4.8 (51)	55.8 (83)	677	21.4(112)	52.6 (46)	76	* *	108	20.4(16)	816	1.1 (20)	816	0.6(8)
Sullivan	382	3.7 (13)	57.6 (107)	356	18.8 (69)	54.8 (74)	61	* *	164	37.8(78)	419	0.2 (5)	419	1.9(22)
Switz""Land	117	6.0 (88)	51.3 (27)	111	12.6 (3)	64.0(123)	17	* *	33	* *	141	0.0 (1)	141	0.7(10)
Tippecanoe	2,887	4.4 (35)	53.8 (62)	2,722	19.1 (79)	53.5 (56)	459	33.8 (22)	390	23.1(28)	3,392	2.2 (46)	3,392	1.6(16)
Tipton Co	127	8.7 (121)	55.9 (86)	119	24.4(122)	47.1 (5)	35	* *	41	* *	175	1.7 (32)	175	8.0(115)
Union Co	107	2.8 (2)	57.9 (109)	100	24.0(121)	49.0 (12)	21	* *	36	* *	136	3.7 (81)	136	2.2(33)
United Fai	978	5.6 (77)	51.7 (32)	924	21.1(111)	51.9 (36)	179	40.8 (46)	246	23.6(33)	1,202	4.8(110)	1,202	6.4(103)
Valpo	826	3.8 (16)	51.7 (32)	782	17.6 (39)	52.8 (48)	173	28.9 (9)	205	21.0(19)	1,092	3.2 (71)	1,092	2.0(26)
Vigo	1,730	5.4 (72)	52.8 (48)	1,459	16.2 (13)	56.8(104)	310	33.2 (21)	1,045	26.8(46)	1,810	2.3 (47)	1,810	2.5(43)
Vincennes	531	7.0 (110)	53.3 (57)	448	16.7 (23)	56.7(103)	83	* *	332	20.2(15)	568	3.5 (75)	568	4.0(67)
W 21st Ave	989	4.8 (51)	57.9 (109)	931	20.4(100)	53.2 (53)	233	64.4 (63)	89	* *	1,182	1.2 (22)	1,182	7.8(113)
Wabash	362	6.1 (92)	52.5 (44)	331	18.7 (65)	51.1 (24)	74	* *	290	19.0(13)	423	4.5(102)	423	6.6(106)
Warren Co	144	4.2 (25)	47.9 (4)	139	19.4 (86)	55.4 (80)	20	* *	57	* *	180	5.6(119)	180	2.2(33)
Warrick Co	605	6.3 (97)	52.9 (49)	569	14.9 (8)	57.3(107)	120	29.2 (11)	136	27.2(50)	744	3.1 (66)	744	5.5(91)
Washington	463	3.7 (13)	56.2 (90)	446	20.6(103)	49.3 (13)	52	* *	56	* *	530	4.0 (92)	530	7.0(110)
Wells Co	353	3.7 (13)	51.6 (31)	336	19.0 (72)	54.2 (65)	75	* *	74	* *	449	1.6 (29)	449	5.1(81)

- (1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Maternal Health Indicators by Clinic
Combined 3 Years

Comparison of Maternal Health Indicators ^(2, 3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Westside	1,526	4.7 (47)	56.6 (93)	1,348	19.9 (97)	51.6 (31)	293	39.6 (41)	389	26.7 (44)	2,141	4.7(107)	2,141	5.0(79)
White Co	546	4.8 (51)	50.5 (18)	529	20.6(103)	52.2 (39)	74	* *	89	* *	650	0.6 (12)	650	0.3(5)
Whitley Co	476	5.9 (86)	50.6 (19)	455	20.2 (99)	52.7 (47)	89	* *	131	21.4(23)	585	3.6 (79)	585	3.2(56)
Wishard	471	3.2 (8)	63.1 (123)	459	17.0 (28)	51.4 (28)	105	35.2 (29)	4	* *	650	7.8(125)	650	6.8(108)
Women""s	1	* *	* *	1	* *	* *	0	* *	0	* *	2	* *	2	* *
Unknown Clinics	295	8.8 (122)	48.5 (6)	1	* *	* *	78	* *	3	* *	4	* *	4	* *
Indiana	109,475	5.0	54.7	101,695	18.7	53.5	21,025	37.0	27,836	26.9	135,164	3.0	135,164	4.1

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6A
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Maternal Behavioral Indicators by Clinic
Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2, 3)										
Clinic	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Action Ctr	59	* *	* *	* *	90	* *	90	* *	45	* *
Albion	455	72.7 (21)	20.7 (26)		654	39.1 (71)	650	14.2 (75)	376	89.1 (46)
Anderson	1,526	60.9 (83)	23.8 (44)		1,935	36.3 (90)	1,935	15.2 (64)	1,022	89.3 (42)
Ball Mem	10	* *	* *	* *	12	* *	12	* *	2	* *
Bartholome	1,066	56.0 (98)	35.4 (88)		1,299	36.0 (92)	1,291	18.4 (31)	661	90.3 (30)
Bedford	555	56.4 (94)	39.3 (99)		749	53.8 (12)	744	13.4 (85)	360	82.2 (96)
Benton Co	77	* *	* *	* *	93	* *	96	* *	52	* *
Berne	86	* *	* *	* *	93	* *	94	* *	52	* *
Bicknell	44	* *	* *	* *	76	* *	75	* *	42	* *
Blackburn	700	70.7 (33)	19.9 (19)		953	34.8 (100)	952	15.9 (56)	506	87.2 (67)
Blackford	185	65.9 (63)	26.5 (58)		240	42.9 (55)	239	18.4 (31)	116	90.5 (28)
Boone Co	364	63.2 (74)	31.3 (76)		468	44.4 (43)	465	11.8 (106)	227	91.2 (25)
Brazil	368	56.3 (95)	39.1 (97)		538	55.4 (6)	536	17.4 (42)	279	86.4 (72)
Brown Co	130	56.2 (96)	39.2 (98)		168	42.3 (58)	168	28.6 (3)	79	* *
Cambridge	54	* *	* *	* *	80	* *	79	* *	34	* *
Carroll Co	144	67.4 (54)	25.7 (53)		201	41.8 (59)	199	16.1 (53)	95	* *
Cass Co	685	59.7 (89)	33.9 (85)		918	43.8 (47)	919	13.2 (88)	478	78.5 (105)
Citizens	704	67.6 (51)	23.7 (41)		1,107	32.1 (111)	1,112	11.8 (106)	564	88.3 (54)
Civic Ctr	1,515	70.1 (37)	23.6 (40)		2,332	37.3 (80)	2,300	11.7 (108)	1,312	91.2 (25)
Clark Co	1,358	62.2 (77)	31.9 (78)		1,793	40.2 (65)	1,786	17.9 (37)	890	91.3 (24)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6A
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Maternal Behavioral Indicators by Clinic
Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2, 3)										
Clinic	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Clay City	23	* *	* *	* *	33	* *	33	* *	10	* *
Clinton	168	72.0 (26)	25.6 (52)		253	37.2 (81)	249	20.5 (15)	123	91.9 (18)
Clinton Co	579	58.7 (92)	32.3 (79)		750	37.2 (81)	754	18.8 (26)	359	89.4 (41)
Comm East	162	71.6 (30)	20.4 (22)		203	34.0(105)	195	7.7 (121)	109	82.6 (95)
Cottage Gr	1,356	73.8 (16)	18.6 (15)		1,842	31.9(112)	1,850	9.5 (115)	989	88.9 (50)
Crawford	122	61.5 (81)	27.0 (59)		176	48.9 (27)	175	15.4 (61)	92	* *
Crown Pt	888	72.7 (21)	19.9 (19)		1,261	34.4(103)	1,249	26.3 (5)	637	93.1 (9)
Daviess Co	276	78.3 (7)	14.9 (3)		447	41.4 (61)	443	12.6 (96)	241	80.9 (100)
Dearborn	476	71.8 (27)	23.1 (36)		764	40.2 (65)	755	15.8 (59)	391	90.3 (30)
Decatur	323	41.2(115)	56.7 (117)		406	53.9 (10)	404	23.5 (8)	212	91.5 (21)
Decatur Co	342	60.8 (84)	35.4 (88)		443	46.3 (31)	432	14.1 (77)	224	81.3 (99)
Dekalb Co	560	67.9 (49)	28.6 (65)		787	43.2 (51)	790	13.9 (79)	393	90.6 (27)
Dubois Co	399	78.4 (6)	17.0 (10)		596	34.2(104)	589	21.6 (14)	293	86.3 (75)
E Chicago	1,054	68.9 (44)	22.5 (33)		1,443	34.7(101)	948	16.1 (53)	670	88.1 (57)
Eaton	18	* *	* *	* *	33	* *	31	* *	12	* *
Elkhart	2,923	40.6(116)	54.8 (115)		3,431	53.9 (10)	3,445	23.2 (9)	1,747	81.9 (97)
Elwood	172	62.8 (76)	27.9 (62)		238	37.4 (78)	237	9.7 (114)	105	91.4 (22)
Fayette Co	320	73.8 (16)	21.3 (29)		521	49.5 (23)	525	12.2 (102)	237	84.0 (90)
Floyd Co	787	60.5 (88)	33.2 (83)		1,116	39.3 (70)	1,096	19.5 (23)	568	89.1 (46)
Forest Man	1,127	75.2 (11)	16.5 (9)		1,632	33.3(109)	1,633	10.0 (112)	852	87.7 (61)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6A
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Maternal Behavioral Indicators by Clinic
Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2, 3)									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Fountain	234	47.0 (110)	46.6 (109)	290	45.2 (36)	286	16.1 (53)	141	85.1 (82)
Franklin C	168	70.8 (31)	26.2 (55)	252	40.5 (64)	249	13.3 (86)	126	89.7 (39)
Fulton Co	293	67.6 (51)	27.6 (61)	426	43.0 (53)	423	14.9 (67)	210	89.0 (49)
Georgetown	2,218	69.8 (40)	20.6 (25)	3,031	29.3 (119)	3,019	9.4 (116)	1,566	88.6 (52)
Gibson Co	296	75.0 (12)	22.0 (31)	449	44.3 (44)	439	14.6 (68)	234	93.6 (6)
Glenn Pk	596	70.1 (37)	21.1 (28)	1,140	31.9 (112)	1,133	12.0 (104)	568	87.0 (69)
Goshen	1,449	45.5 (113)	50.7 (113)	1,694	52.7 (16)	1,690	12.9 (92)	820	80.9 (100)
Grant Co	930	73.1 (20)	20.5 (24)	1,420	39.8 (69)	1,422	12.8 (94)	693	90.0 (33)
Grant St	597	73.4 (19)	14.9 (3)	936	32.7 (110)	927	8.2 (120)	451	83.8 (91)
Grassy Crk	1,522	67.0 (57)	26.4 (57)	2,172	35.7 (96)	2,161	12.0 (104)	1,093	85.6 (79)
Greene Co	238	72.7 (21)	20.2 (21)	418	44.7 (41)	320	16.9 (47)	190	90.0 (33)
Hamilton	1,075	71.8 (27)	22.7 (34)	1,556	34.6 (102)	1,541	20.2 (18)	803	91.9 (18)
Hammond	1,122	69.7 (41)	20.9 (27)	1,669	31.9 (112)	1,637	13.8 (83)	831	87.4 (64)
Hancock Co	460	65.4 (67)	29.8 (72)	654	36.7 (84)	653	19.8 (20)	349	91.7 (20)
Hanna	2,186	62.1 (79)	30.2 (75)	2,846	34.0 (105)	2,838	13.1 (89)	1,597	82.8 (94)
Harrison	371	56.1 (97)	36.9 (95)	567	50.3 (22)	566	20.3 (16)	288	95.1 (2)
Hendricks	596	78.7 (4)	15.1 (6)	953	30.3 (118)	935	22.4 (12)	498	92.6 (12)
Henry Co	642	67.0 (57)	29.8 (72)	835	49.3 (25)	832	13.9 (79)	402	81.6 (98)
Hessville	618	69.3 (43)	23.5 (39)	884	35.6 (98)	868	19.8 (20)	436	93.1 (9)
Hispanic C	218	64.7 (68)	23.4 (37)	287	31.7 (115)	288	22.6 (11)	159	89.9 (37)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6A
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Maternal Behavioral Indicators by Clinic
Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2, 3)										
Clinic	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	% (Rank)		Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Howard Co	1,347	60.7 (85)	32.4 (81)		1,769	43.1 (52)	1,762	10.5 (110)	892	86.4 (72)
Huntington	477	40.5(117)	56.4 (116)		624	63.9 (2)	624	18.3 (34)	303	84.2 (87)
IN0108 Closed	42	* *	* *		51	* *	50	* *	0	* *
IN0904 Closed	184	79.3 (3)	15.8 (8)		289	39.1 (71)	286	10.5 (110)	2	* *
IN2224 Closed	1	* *	* *		1	* *	1	* *	1	* *
IN2305 Closed	6	* *	* *		14	* *	14	* *	0	* *
Jackson Co	670	70.0 (39)	23.7 (41)		905	44.2 (45)	897	14.6 (68)	457	92.1 (16)
Jasonville	13	* *	* *		21	* *	20	* *	9	* *
Jasper Co	390	51.3(104)	43.6 (104)		534	53.7 (13)	535	16.4 (49)	259	84.6 (84)
Jay Co	266	72.2 (25)	23.7 (41)		368	49.2 (26)	364	14.3 (72)	176	90.3 (30)
Jefferson	480	60.6 (86)	36.3 (93)		589	46.9 (29)	583	14.4 (71)	288	86.1 (77)
Jennings	624	28.0(118)	68.3 (118)		756	65.7 (1)	756	12.3 (101)	362	77.9 (106)
Johnson Co	1,303	66.2 (62)	26.3 (56)		1,636	31.7(115)	1,610	17.2 (44)	795	89.6 (40)
Kosciusko	732	76.8 (9)	14.9 (3)		1,005	29.3(119)	1,002	16.4 (49)	542	88.0 (58)
Lagrange	227	79.7 (2)	12.8 (2)		342	36.0 (92)	337	16.3 (51)	182	87.4 (64)
Lake Station	633	64.5 (70)	28.9 (67)		1,012	41.0 (63)	1,002	14.3 (72)	504	84.5 (85)
Laporte	895	54.3 (99)	37.5 (96)		1,139	41.2 (62)	1,126	18.3 (34)	541	94.6 (3)
Main St	1,361	72.4 (24)	22.7 (34)		1,948	43.7 (49)	1,926	17.8 (38)	1,005	90.4 (29)
Marshall	688	65.7 (65)	28.6 (65)		860	36.5 (86)	857	18.4 (31)	444	87.6 (62)
Martin Co	94	* *	* *		142	54.9 (8)	143	21.7 (13)	72	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6A
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Maternal Behavioral Indicators by Clinic
Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2, 3)										
Clinic	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	% (Rank)		Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Memorial	340	67.4 (54)	25.0 (49)		454	35.7 (96)	450	14.2 (75)	223	93.3 (7)
Merrillville	0	* *	* *		3	* *	3	* *	1	* *
Methodist	413	71.7 (29)	17.9 (11)		562	37.4 (78)	561	8.4 (119)	257	82.9 (93)
Miami Co	370	68.4 (46)	29.2 (68)		561	48.1 (28)	556	18.0 (36)	273	87.9 (59)
Mich City	723	64.0 (71)	27.5 (60)		1,020	36.5 (86)	763	23.9 (7)	508	92.7 (11)
Moms/Kids	126	68.3 (47)	25.4 (51)		183	36.6 (85)	183	17.5 (41)	120	89.2 (43)
Monroe Co	1,142	63.3 (73)	32.3 (79)		1,635	46.1 (33)	1,088	29.1 (2)	779	91.4 (22)
Montgomery	621	46.1(112)	48.0 (110)		719	44.9 (37)	719	15.9 (56)	344	89.2 (43)
Morgan Co	691	64.7 (68)	29.5 (70)		933	36.4 (89)	927	18.8 (26)	461	90.0 (33)
N Arlingto	1,397	75.0 (12)	18.2 (13)		1,812	36.3 (90)	1,603	14.0 (78)	957	86.3 (75)
N Liberty	31	* *	* *		40	* *	41	* *	21	* *
N Manchest	108	65.7 (65)	32.4 (81)		151	43.0 (53)	148	18.9 (24)	70	* *
Newton Co	108	67.6 (51)	24.1 (46)		138	39.9 (68)	141	11.3 (109)	72	* *
Northwest	1,019	70.8 (31)	19.1 (17)		1,339	24.6(123)	1,334	12.7 (95)	660	92.4 (13)
Ohio Co	14	* *	* *		22	* *	22	* *	9	* *
Orange Co	310	52.9(101)	45.5 (106)		409	57.0 (5)	410	17.8 (38)	205	84.4 (86)
Owen Co	242	51.2(105)	46.3 (108)		333	51.7 (18)	331	13.9 (79)	156	76.9 (107)
Parke Co	152	52.6(102)	42.8 (102)		198	50.5 (21)	178	23.0 (10)	89	* *
Perry Co	238	62.2 (77)	33.2 (83)		294	42.5 (56)	291	2.4 (124)	160	94.4 (4)
Pike Co	99	* *	* *		177	41.8 (59)	175	12.6 (96)	93	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6A
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Maternal Behavioral Indicators by Clinic
Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2, 3)										
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption				
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)		
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	
Portage	607	66.4 (61)	28.5 (64)	925	43.7 (49)	916	19.8 (20)	484	88.2 (55)	
Posey Co	162	66.7 (59)	29.6 (71)	229	45.4 (35)	226	15.9 (56)	118	87.3 (66)	
Pulaski Co	129	65.9 (63)	27.9 (62)	198	51.5 (19)	198	18.7 (29)	85	* *	
Purdue	280	52.1(103)	43.6 (104)	352	55.1 (7)	220	64.1 (1)	157	95.5 (1)	
Putnam Co	463	46.4(111)	50.1 (112)	604	54.0 (9)	606	17.0 (45)	291	85.6 (79)	
Randolph	323	83.3 (1)	11.1 (1)	506	35.8 (95)	500	15.8 (59)	256	92.2 (14)	
Richmond	1,151	50.5(106)	42.8 (102)	1,450	53.5 (15)	1,364	13.1 (89)	700	74.4 (109)	
Ripley Co	344	63.7 (72)	31.4 (77)	468	40.0 (67)	470	13.0 (91)	239	84.1 (89)	
Rush Co	165	69.7 (41)	26.1 (54)	252	51.2 (20)	252	9.1 (118)	127	75.6 (108)	
SB Ireland Rd	1,420	75.0 (12)	20.4 (22)	1,872	38.1 (75)	1,856	18.9 (24)	919	93.3 (7)	
SB Lafayette	1,665	68.7 (45)	24.0 (45)	2,229	37.6 (77)	2,215	14.3 (72)	1,181	89.2 (43)	
Scott Co	495	59.2 (91)	36.6 (94)	596	53.7 (13)	598	15.1 (65)	296	83.1 (92)	
Shelby Co	492	67.9 (49)	24.8 (48)	705	36.5 (86)	709	12.1 (103)	380	87.1 (68)	
Shelters	9	* *	* *	17	* *	17	* *	8	* *	
SJ Reg Ctr	112	78.6 (5)	15.2 (7)	147	35.4 (99)	146	17.8 (38)	65	* *	
So Calhoun	4,559	57.5 (93)	35.9 (91)	6,081	44.8 (39)	6,065	13.9 (79)	3,043	88.7 (51)	
Spencer Co	171	66.7 (59)	29.2 (68)	220	35.9 (94)	223	7.2 (122)	126	85.7 (78)	
St Francis	0	* *	* *	0	* *	0	* *	0	* *	
St V-FLC	290	73.8 (16)	19.0 (16)	371	26.7(122)	369	13.3 (86)	169	86.4 (72)	
Starke Co	370	53.5(100)	41.9 (101)	463	52.7 (16)	467	15.4 (61)	230	80.9 (100)	

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6A
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Maternal Behavioral Indicators by Clinic
Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2, 3)										
Clinic	Medical Care			WIC Enrollment			Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)		
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	
Steuben Co	517	47.2(108)	49.9 (111)	709	62.2 (3)	701	12.6 (96)	366	80.6 (103)	
Sullivan	196	74.5 (15)	22.4 (32)	381	46.2 (32)	381	16.3 (51)	203	85.2 (81)	
Switz""Land	90	* *	* *	118	39.0 (73)	117	15.4 (61)	51	* *	
Tippecanoe	2,410	47.1(109)	46.0 (107)	2,886	49.5 (23)	2,789	20.3 (16)	1,417	85.0 (83)	
Tipton Co	91	* *	* *	129	38.0 (76)	128	18.8 (26)	78	* *	
Union Co	72	* *	* *	109	38.5 (74)	107	18.7 (29)	50	* *	
United Fai	744	59.3 (90)	35.5 (90)	982	43.9 (46)	978	20.1 (19)	462	94.4 (4)	
Valpo	570	70.4 (35)	24.7 (47)	828	37.2 (81)	826	28.0 (4)	424	92.0 (17)	
Vigo	947	76.9 (8)	18.1 (12)	1,744	42.5 (56)	1,732	12.4 (100)	908	88.2 (55)	
Vincennes	324	76.5 (10)	21.3 (29)	541	43.8 (47)	535	6.4 (123)	294	89.1 (46)	
W 21st Ave	706	68.3 (47)	19.7 (18)	984	34.0(105)	990	12.5 (99)	438	79.7 (104)	
Wabash	252	67.1 (56)	29.8 (72)	366	44.8 (39)	364	12.9 (92)	196	89.8 (38)	
Warren Co	123	49.6(107)	41.5 (100)	144	34.0(105)	145	16.6 (48)	76	* *	
Warrick Co	447	60.6 (86)	36.2 (92)	609	44.7 (41)	607	14.5 (70)	323	87.6 (62)	
Washington	370	43.2(114)	51.1 (114)	460	58.5 (4)	463	15.1 (65)	221	86.9 (70)	
Wells Co	258	61.2 (82)	34.9 (87)	361	44.9 (37)	353	17.3 (43)	179	92.2 (14)	
Westside	1,095	63.2 (74)	23.4 (37)	1,529	31.3(117)	1,525	9.9 (113)	722	86.6 (71)	
White Co	395	70.6 (34)	25.3 (50)	550	45.6 (34)	550	13.6 (84)	284	84.2 (87)	
Whitley Co	337	62.0 (80)	34.7 (86)	479	46.6 (30)	475	26.1 (6)	221	90.0 (33)	
Wishard	396	70.2 (36)	18.2 (13)	473	28.5(121)	464	9.3 (117)	148	87.8 (60)	

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6A
2009 - 2011 Pregnancy Nutrition Surveillance (1)
Indiana
Comparison of Maternal Behavioral Indicators by Clinic
Combined 3 Years

Comparison of Maternal Behavioral Indicators (2, 3)											
Clinic	Medical Care					WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care			1st Trimester		Pregnancy (4)		During Pregnancy (5)	
	Number	%	%	%	%	Number	%	Number	%	Number	%
	(Rank)	(Rank)	(Rank)	(Rank)	(Rank)	(Rank)	(Rank)	(Rank)	(Rank)	(Rank)	(Rank)
Women''s	1	*	*	*	*	1	*	*	1	*	*
Unknown Clinics	1	*	*	*	*	305	16.7 (124)	294	17.0 (45)	296	88.5 (53)
Indiana	80,115	63.3	30.0			110,014	40.7	107,563	15.6	55,435	87.7

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2009 - 2011 Pregnancy Nutrition Surveillance (1)
Indiana
Comparison of Smoking Indicators by Clinic
Combined 3 Years

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Action Ctr	120	14.2 (9)	113	8.8 (7)	15	* *	89	* *	111	24.3 (35)
Albion	826	30.8 (43)	788	27.3 (81)	237	22.8 (99)	649	39.0 (46)	781	30.7 (42)
Anderson	2,608	34.5 (66)	2,538	26.0 (72)	864	33.6 (53)	1,933	49.2 (78)	2,515	45.3 (81)
Ball Mem	22	* *	22	* *	7	* *	11	* *	22	* *
Bartholome	1,632	35.8 (74)	1,590	25.3 (68)	564	35.6 (37)	1,289	51.6 (86)	1,570	46.5 (88)
Bedford	931	40.5 (103)	901	27.7 (84)	363	36.6 (29)	745	50.3 (82)	895	46.4 (87)
Benton Co	122	32.8 (53)	118	28.0 (87)	40	* *	96	* *	116	48.3 (92)
Berne	113	27.4 (35)	112	16.1 (25)	31	* *	94	* *	111	35.1 (55)
Bicknell	91	* *	86	* *	35	* *	75	* *	83	* *
Blackburn	1,361	17.8 (15)	1,313	11.7 (13)	234	49.1 (6)	939	23.0 (15)	1,304	18.6 (23)
Blackford	293	42.3 (110)	289	31.8 (112)	123	29.3 (76)	238	47.9 (72)	286	43.7 (77)
Boone Co	612	39.1 (89)	586	29.2 (96)	223	33.2 (54)	464	51.3 (84)	579	48.4 (93)
Brazil	641	40.6 (104)	623	27.8 (86)	251	33.9 (50)	536	56.9 (108)	615	52.5 (110)
Brown Co	209	34.4 (65)	202	19.3 (37)	69	* *	166	50.0 (80)	200	44.0 (78)
Cambridge	105	34.3 (62)	101	19.8 (39)	34	* *	79	* *	99	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2009 - 2011 Pregnancy Nutrition Surveillance (1)
Indiana
Comparison of Smoking Indicators by Clinic
Combined 3 Years

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Carroll Co	248	33.1 (57)	246	28.0 (87)	82	* *	199	31.7 (31)	243	14.8 (13)
Cass Co	1,148	25.3 (30)	1,082	18.4 (32)	269	29.7 (73)	917	27.3 (23)	1,067	16.5 (15)
Citizens	1,852	17.4 (12)	1,635	17.5 (28)	254	37.8 (27)	1,096	35.7 (40)	1,620	33.6 (48)
Civic Ctr	2,754	37.2 (81)	2,421	20.1 (41)	868	41.6 (17)	2,286	25.0 (18)	2,376	8.5 (5)
Clark Co	2,271	34.3 (62)	2,188	23.9 (61)	742	35.0 (40)	1,783	41.4 (56)	2,157	37.1 (62)
Clay City	38	* *	37	* *	9	* *	33	* *	37	* *
Clinton	303	44.6 (117)	297	34.3 (122)	131	28.2 (84)	248	64.5 (122)	294	58.8 (123)
Clinton Co	901	28.0 (38)	882	19.7 (38)	246	34.6 (43)	754	23.2 (16)	870	10.8 (7)
Comm East	330	24.5 (26)	328	23.2 (58)	80	* *	170	39.4 (49)	327	41.9 (71)
Cottage Gr	2,538	30.4 (42)	2,413	30.7 (105)	714	18.8 (103)	1,793	44.2 (63)	2,357	43.2 (76)
Crawford	219	42.9 (113)	212	38.7 (127)	90	* *	175	64.0 (121)	209	52.6 (111)
Crown Pt	1,729	27.4 (35)	1,664	14.1 (22)	451	51.2 (4)	1,244	37.5 (42)	1,637	35.8 (58)
Daviess Co	555	33.9 (61)	514	22.0 (49)	167	34.1 (48)	446	44.4 (64)	502	42.4 (74)
Dearborn	957	45.5 (120)	922	30.8 (107)	412	36.9 (28)	754	53.7 (96)	903	48.7 (95)
Decatur	506	36.2 (76)	487	23.0 (56)	175	36.6 (29)	403	44.7 (68)	479	40.9 (68)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2009 - 2011 Pregnancy Nutrition Surveillance (1)
Indiana
Comparison of Smoking Indicators by Clinic
Combined 3 Years

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Decatur Co	555	39.8 (95)	540	30.4 (103)	212	26.4 (91)	434	60.6 (115)	536	57.3 (120)
Dekalb Co	986	38.3 (86)	953	31.6 (111)	362	25.4 (94)	790	58.7 (112)	943	53.4 (113)
Dubois Co	782	26.6 (33)	767	18.6 (34)	203	34.5 (44)	578	41.0 (53)	757	37.8 (63)
E Chicago	1,852	13.8 (8)	1,805	5.8 (4)	246	56.9 (1)	1,438	19.1 (8)	1,784	13.5 (11)
Eaton	44	* *	42	* *	13	* *	30	* *	42	* *
Elkhart	4,070	25.0 (29)	3,927	15.9 (24)	961	30.4 (66)	3,437	23.7 (17)	3,881	9.3 (6)
Elwood	312	38.5 (88)	305	36.1 (125)	117	20.5 (102)	238	58.8 (113)	299	59.2 (126)
Fayette Co	649	39.9 (96)	631	29.5 (97)	249	36.1 (36)	524	59.2 (114)	626	55.9 (116)
Floyd Co	1,409	38.2 (85)	1,342	25.1 (67)	508	40.0 (18)	1,093	48.9 (77)	1,313	46.0 (85)
Forest Man	2,228	18.9 (17)	2,104	12.7 (16)	389	35.0 (40)	1,067	26.4 (21)	1,518	23.6 (32)
Fountain	356	46.9 (123)	343	31.5 (110)	162	36.4 (33)	285	61.4 (116)	339	57.5 (121)
Franklin C	325	39.1 (89)	316	28.5 (91)	124	37.9 (26)	249	55.0 (101)	310	52.3 (106)
Fulton Co	517	35.6 (72)	491	24.8 (65)	175	36.6 (29)	422	38.9 (45)	486	33.5 (47)
Georgetown	4,277	11.3 (5)	4,077	6.9 (5)	447	51.0 (5)	3,025	18.6 (7)	4,007	17.1 (19)
Gibson Co	577	41.6 (108)	549	34.2 (121)	222	28.8 (81)	441	57.8 (110)	545	54.3 (114)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2009 - 2011 Pregnancy Nutrition Surveillance (1)
Indiana
Comparison of Smoking Indicators by Clinic
Combined 3 Years

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Glenn Pk	1,894	11.0 (4)	1,576	7.2 (6)	149	48.3 (9)	1,133	20.4 (12)	1,559	17.4 (20)
Goshen	2,035	21.0 (19)	1,968	12.5 (15)	415	38.3 (25)	1,685	18.0 (5)	1,951	4.7 (1)
Grant Co	1,756	34.5 (66)	1,684	28.2 (90)	572	25.5 (92)	1,424	42.6 (60)	1,667	23.9 (34)
Grant St	1,207	17.4 (12)	1,171	12.9 (17)	202	36.6 (29)	929	25.8 (20)	1,151	24.7 (36)
Grassy Crk	3,216	17.7 (14)	2,978	10.7 (11)	502	48.4 (8)	2,150	25.3 (19)	2,945	22.3 (27)
Greene Co	521	39.3 (93)	509	28.5 (91)	196	29.1 (77)	387	55.8 (104)	506	45.7 (83)
Hamilton	2,045	22.2 (22)	1,984	14.0 (21)	435	39.8 (19)	1,539	29.9 (27)	1,959	23.3 (31)
Hammond	2,387	15.2 (10)	2,199	9.3 (9)	323	51.7 (3)	1,643	26.4 (21)	2,171	22.1 (26)
Hancock Co	866	34.5 (66)	834	22.7 (54)	289	34.3 (47)	650	34.5 (36)	823	16.8 (17)
Hanna	3,992	21.6 (20)	3,813	18.0 (30)	793	32.4 (57)	2,822	32.7 (33)	3,748	31.4 (45)
Harrison	727	40.0 (97)	694	26.5 (74)	278	39.2 (21)	565	52.2 (89)	682	45.3 (81)
Hendricks	1,273	31.3 (46)	1,136	15.7 (23)	324	46.3 (11)	943	43.3 (62)	1,114	39.4 (66)
Henry Co	1,003	46.1 (121)	979	31.8 (112)	447	30.6 (64)	827	53.0 (92)	966	49.5 (97)
Hessville	1,141	22.3 (23)	1,108	10.8 (12)	239	44.8 (13)	869	20.0 (10)	1,097	15.0 (14)
Hispanic C	370	4.6 (2)	351	0.9 (1)	15	* *	288	14.6 (3)	339	8.3 (4)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2009 - 2011 Pregnancy Nutrition Surveillance (1)
Indiana
Comparison of Smoking Indicators by Clinic
Combined 3 Years

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Howard Co	2,164	35.2 (71)	2,106	27.0 (79)	738	25.5 (92)	1,762	44.6 (66)	2,076	36.6 (61)
Huntington	726	49.9 (127)	702	31.2 (109)	344	30.8 (63)	621	50.6 (83)	694	46.3 (86)
IN0108 Closed	60	* *	58	* *	12	* *	51	* *	58	* *
IN0904 Closed	323	50.5 (128)	269	26.0 (72)	127	42.5 (16)	283	40.6 (50)	264	17.0 (18)
IN2224 Closed	1	* *	1	* *	0	* *	1	* *	1	* *
IN2305 Closed	14	* *	9	* *	3	* *	13	* *	8	* *
Jackson Co	1,070	36.3 (77)	1,046	22.9 (55)	375	38.7 (22)	895	39.1 (47)	1,040	34.6 (54)
Jasonville	24	* *	24	* *	7	* *	21	* *	24	* *
Jasper Co	697	32.6 (52)	652	27.5 (83)	209	21.5 (101)	539	57.5 (109)	648	52.3 (106)
Jay Co	465	35.1 (70)	450	25.8 (71)	156	29.5 (75)	360	42.5 (59)	440	36.4 (60)
Jefferson	733	42.0 (109)	704	34.7 (124)	293	27.0 (87)	581	40.6 (50)	694	28.4 (41)
Jennings	864	37.4 (82)	840	30.6 (104)	310	15.5 (104)	756	56.0 (105)	829	50.9 (101)
Johnson Co	2,227	30.8 (43)	2,156	23.1 (57)	649	32.5 (56)	1,624	41.9 (58)	2,116	34.2 (51)
Kosciusko	1,275	30.8 (43)	1,205	22.0 (49)	356	32.3 (58)	1,000	44.4 (64)	1,190	34.1 (50)
Lagrange	434	31.6 (48)	412	23.8 (59)	125	30.4 (66)	337	48.4 (76)	400	42.5 (75)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2009 - 2011 Pregnancy Nutrition Surveillance (1)
Indiana
Comparison of Smoking Indicators by Clinic
Combined 3 Years

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Lake Station	1,585	22.5 (24)	1,399	18.9 (36)	294	38.4 (24)	1,001	42.7 (61)	1,385	40.1 (67)
Laporte	1,381	37.8 (83)	1,304	27.4 (82)	477	30.0 (70)	1,132	52.7 (90)	1,279	50.9 (101)
Main St	2,554	34.6 (69)	2,451	28.0 (87)	842	28.5 (82)	1,851	35.3 (39)	2,422	23.2 (30)
Marshall	1,063	32.9 (55)	1,029	21.9 (48)	331	34.4 (46)	857	20.1 (11)	1,020	5.2 (2)
Martin Co	172	43.0 (114)	166	30.7 (105)	68	* *	143	41.3 (55)	165	25.5 (37)
Memorial	534	24.5 (26)	531	13.4 (18)	130	43.1 (15)	450	16.4 (4)	531	14.1 (12)
Merrillville	8	* *	5	* *	0	* *	3	* *	5	* *
Methodist	812	23.0 (25)	806	22.0 (49)	186	30.6 (64)	518	37.1 (41)	804	40.9 (68)
Miami Co	670	40.0 (97)	637	24.5 (63)	247	39.3 (20)	555	54.4 (99)	634	45.9 (84)
Mich City	1,353	25.4 (31)	1,263	21.4 (45)	313	34.5 (44)	1,014	32.3 (32)	1,247	27.5 (40)
Moms/Kids	253	17.0 (11)	231	9.1 (8)	38	* *	183	28.4 (25)	228	20.2 (25)
Monroe Co	2,023	34.3 (62)	1,919	21.2 (44)	644	33.7 (51)	1,506	30.6 (29)	1,889	17.5 (21)
Montgomery	872	41.5 (107)	843	31.1 (108)	350	28.9 (80)	718	56.4 (107)	834	49.2 (96)
Morgan Co	1,275	38.4 (87)	1,228	33.6 (120)	462	26.6 (89)	917	56.3 (106)	1,214	55.0 (115)
N Arlingto	2,448	22.0 (21)	2,335	16.8 (26)	488	38.7 (22)	1,492	34.2 (35)	2,278	34.2 (51)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2009 - 2011 Pregnancy Nutrition Surveillance (1)
Indiana
Comparison of Smoking Indicators by Clinic
Combined 3 Years

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
N Liberty	51	* *	50	* *	18	* *	40	* *	50	* *
N Manchest	181	33.1 (57)	172	20.3 (42)	56	* *	148	35.1 (38)	171	25.7 (38)
Newton Co	185	27.6 (37)	180	30.0 (100)	47	* *	141	66.7 (123)	180	58.9 (124)
Northwest	1,959	7.8 (3)	1,877	4.3 (3)	141	56.7 (2)	1,323	12.2 (2)	1,862	11.9 (8)
Ohio Co	29	* *	29	* *	10	* *	22	* *	27	* *
Orange Co	487	42.3 (110)	471	29.5 (97)	195	26.7 (88)	411	41.6 (57)	467	19.5 (24)
Owen Co	388	47.2 (124)	377	32.4 (115)	177	28.2 (84)	333	58.0 (111)	372	50.5 (100)
Parke Co	240	44.2 (115)	240	28.8 (94)	105	35.2 (39)	198	54.0 (98)	237	51.9 (105)
Perry Co	386	44.6 (117)	374	38.0 (126)	161	24.8 (96)	289	61.9 (118)	368	59.0 (125)
Pike Co	226	31.9 (49)	216	24.1 (62)	71	* *	174	63.2 (119)	214	57.0 (119)
Portage	1,193	33.5 (59)	1,141	20.4 (43)	378	46.3 (11)	914	44.6 (66)	1,121	42.0 (72)
Posey Co	293	46.4 (122)	275	33.1 (118)	125	29.6 (74)	226	67.3 (124)	273	56.4 (117)
Pulaski Co	264	33.7 (60)	243	32.5 (116)	81	* *	197	51.3 (84)	240	45.0 (80)
Purdue	427	2.8 (1)	413	1.7 (2)	11	* *	351	6.8 (1)	405	5.2 (2)
Putnam Co	732	40.8 (106)	706	26.9 (78)	285	35.4 (38)	601	52.1 (88)	700	48.1 (90)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2009 - 2011 Pregnancy Nutrition Surveillance (1)
Indiana
Comparison of Smoking Indicators by Clinic
Combined 3 Years

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Randolph	622	33.0 (56)	605	25.6 (70)	196	25.0 (95)	498	45.0 (69)	584	37.8 (63)
Richmond	1,733	36.5 (78)	1,682	25.5 (69)	609	29.1 (77)	1,445	48.0 (74)	1,661	39.1 (65)
Ripley Co	610	36.9 (80)	582	32.3 (114)	209	24.4 (97)	468	53.2 (93)	569	52.4 (109)
Rush Co	312	40.1 (100)	305	29.8 (99)	120	30.0 (70)	253	61.7 (117)	302	56.6 (118)
SB Ireland Rd	2,385	32.1 (51)	2,275	17.1 (27)	713	48.9 (7)	1,849	30.0 (28)	2,253	22.5 (28)
SB Lafayette	2,786	24.8 (28)	2,676	13.9 (20)	653	47.2 (10)	2,216	22.1 (13)	2,625	13.1 (10)
Scott Co	720	45.3 (119)	697	34.6 (123)	309	27.2 (86)	595	55.6 (103)	690	50.4 (99)
Shelby Co	929	36.1 (75)	894	26.8 (77)	310	31.3 (62)	706	52.7 (90)	885	51.3 (103)
Shelters	24	* *	21	* *	5	* *	16	* *	21	* *
SJ Reg Ctr	178	30.3 (40)	178	11.8 (14)	54	* *	146	19.2 (9)	178	18.0 (22)
So Calhoun	7,402	26.4 (32)	7,178	17.6 (29)	1,855	33.7 (51)	6,029	34.5 (36)	7,075	31.9 (46)
Spencer Co	285	36.5 (78)	281	23.8 (59)	103	34.0 (49)	222	46.8 (71)	274	48.2 (91)
St Francis	10	* *	10	* *	0	* *	0	* *	10	* *
St V-FLC	517	12.0 (6)	504	9.5 (10)	61	* *	346	18.5 (6)	504	26.2 (39)
Starke Co	566	40.1 (100)	550	27.1 (80)	217	32.7 (55)	464	53.9 (97)	547	51.4 (104)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2009 - 2011 Pregnancy Nutrition Surveillance (1)
Indiana
Comparison of Smoking Indicators by Clinic
Combined 3 Years

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Steuben Co	838	40.0 (97)	816	26.5 (74)	324	30.2 (68)	701	54.8 (100)	802	49.8 (98)
Sullivan	442	42.8 (112)	419	33.4 (119)	172	22.7 (100)	381	53.5 (95)	417	47.5 (89)
Switz""Land	148	49.3 (126)	141	32.6 (117)	68	* *	118	53.4 (94)	138	48.6 (94)
Tippecanoe	3,521	27.0 (34)	3,392	19.9 (40)	906	26.6 (89)	2,829	38.2 (43)	3,349	33.9 (49)
Tipton Co	182	31.9 (49)	175	30.3 (102)	54	* *	129	49.6 (79)	174	35.6 (57)
Union Co	143	31.5 (47)	136	25.0 (66)	42	* *	107	55.1 (102)	132	52.3 (106)
United Fai	1,241	30.3 (40)	1,202	21.5 (46)	364	29.1 (77)	976	39.2 (48)	1,188	35.2 (56)
Valpo	1,131	32.8 (53)	1,092	18.8 (35)	358	44.7 (14)	822	28.7 (26)	1,069	16.7 (16)
Vigo	2,069	39.4 (94)	1,810	24.6 (64)	678	36.3 (34)	1,734	46.3 (70)	1,794	42.1 (73)
Vincennes	649	37.8 (83)	568	21.8 (47)	212	32.1 (60)	533	41.1 (54)	551	35.9 (59)
W 21st Ave	1,229	19.9 (18)	1,182	18.5 (33)	234	36.3 (34)	987	32.8 (34)	1,173	31.2 (44)
Wabash	453	40.2 (102)	423	30.0 (100)	165	31.5 (61)	362	40.6 (50)	415	23.1 (29)
Warren Co	185	48.1 (125)	180	26.7 (76)	85	* *	144	63.9 (120)	180	58.3 (122)
Warrick Co	777	40.7 (105)	744	27.7 (84)	302	30.1 (69)	607	51.7 (87)	737	53.2 (112)
Washington	544	44.3 (116)	530	29.1 (95)	236	32.2 (59)	463	47.9 (72)	521	34.4 (53)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2009 - 2011 Pregnancy Nutrition Surveillance (1)
Indiana
Comparison of Smoking Indicators by Clinic
Combined 3 Years

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Wells Co	460	39.1 (89)	449	28.7 (93)	174	29.9 (72)	354	48.3 (75)	444	41.4 (70)
Westside	2,314	17.9 (16)	2,141	18.2 (31)	357	28.3 (83)	1,518	31.5 (30)	2,109	30.8 (43)
White Co	668	30.1 (39)	650	22.2 (52)	191	23.0 (98)	547	27.6 (24)	641	12.5 (9)
Whitley Co	604	35.6 (72)	585	22.2 (52)	206	35.0 (40)	474	50.2 (81)	583	44.4 (79)
Wishard	655	13.6 (7)	650	13.4 (18)	88	* *	433	22.4 (14)	639	23.6 (32)
Women""s	2	* *	2	* *	0	* *	1	* *	2	* *
Unknown Clinics	307	39.1 (89)	4	* *	1	* *	291	38.5 (44)	3	* *
Indiana	142,169	29.0	135,160	20.5	38,415	34.2	107,925	37.5	132,772	31.6

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Infant Health Indicators by Clinic
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)									
Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)						
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Action Ctr	111	7.2 (41)	4.5 (15)	81	* *	72	* *	111	62.2 (74)
Albion	762	7.7 (67)	6.3 (57)	605	16.2 (101)	506	4.0(110)	763	74.0 (17)
Anderson	2,485	8.2 (79)	5.6 (41)	1,849	14.1 (75)	1,580	3.3 (90)	2,487	58.6 (96)
Ball Mem	22	* *	* *	12	* *	11	* *	22	* *
Bartholome	1,542	6.8 (30)	6.5 (62)	1,242	13.8 (65)	1,069	2.1 (33)	1,542	68.4 (45)
Bedford	872	9.7(106)	5.3 (35)	703	10.4 (10)	627	4.6(121)	874	65.4 (62)
Benton Co	112	9.8(108)	7.1 (87)	87	* *	69	* *	112	64.3 (69)
Berne	109	9.2 (97)	4.6 (18)	90	* *	77	* *	109	82.6 (4)
Bicknell	76	* *	* *	65	* *	54	* *	76	* *
Blackburn	1,300	10.1(116)	5.2 (33)	898	14.5 (81)	768	3.4 (93)	1,300	52.8(116)
Blackford	277	7.6 (59)	5.8 (49)	229	11.8 (23)	202	1.5 (15)	277	69.7 (41)
Boone Co	568	8.1 (76)	6.2 (56)	435	12.2 (31)	382	2.9 (67)	568	56.7(104)
Brazil	597	4.7 (3)	7.4 (93)	503	10.1 (6)	452	1.1 (8)	596	61.1 (80)
Brown Co	192	6.3 (19)	10.9 (126)	158	12.7 (42)	137	0.0 (1)	194	80.9 (5)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Infant Health Indicators by Clinic
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)										
Clinic	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)		Number	%	Number	%	Number	%
	Number	% (Rank)	% (Rank)	(Rank)						
Cambridge	97	* * * *	* * * *	74	* *	66	* *	97	* *	
Carroll Co	236	8.5 (85)	9.3 (123)	190	14.7 (84)	162	3.1 (81)	236	72.9 (23)	
Cass Co	1,032	6.6 (24)	6.5 (62)	838	9.8 (5)	748	2.4 (40)	1,038	60.9 (82)	
Citizens	1,586	9.6(103)	5.2 (33)	875	13.9 (69)	747	3.6(100)	1,593	55.0(108)	
Civic Ctr	2,337	9.4(100)	5.5 (39)	1,976	16.1 (100)	1,643	2.9 (67)	2,343	59.8 (88)	
Clark Co	2,099	8.4 (83)	5.6 (41)	1,672	13.6 (59)	1,437	3.7(104)	2,097	67.8 (48)	
Clay City	34	* * * *	* * * *	29	* *	26	* *	34	* *	
Clinton	286	8.0 (74)	5.9 (50)	243	14.4 (79)	207	2.4 (40)	286	54.5(111)	
Clinton Co	853	6.2 (15)	9.3 (123)	721	12.2 (31)	632	1.9 (26)	852	72.4 (26)	
Comm East	320	11.3(123)	3.8 (3)	199	19.1 (119)	160	5.0(122)	322	39.8(126)	
Cottage Gr	2,317	9.1 (96)	7.0 (84)	1,682	14.3 (77)	1,427	2.7 (59)	2,318	57.2(101)	
Crawford	202	7.4 (51)	5.4 (37)	164	14.0 (71)	141	1.4 (11)	201	67.7 (50)	
Crown Pt	1,624	6.2 (15)	7.8 (101)	1,180	11.9 (25)	1,038	2.2 (36)	1,626	85.4 (3)	
Daviess Co	479	7.1 (37)	2.9 (1)	394	15.7 (96)	328	3.0 (71)	485	58.4 (97)	

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Infant Health Indicators by Clinic
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)									
Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)						
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Dearborn	880	7.5 (57)	8.9 (116)	718	11.8 (23)	624	2.4 (40)	892	61.1 (80)
Decatur	463	7.3 (44)	5.4 (37)	377	11.7 (21)	333	3.0 (71)	463	78.0 (9)
Decatur Co	528	7.6 (59)	5.7 (44)	420	13.8 (65)	362	3.3 (90)	528	66.5 (56)
Dekalb Co	904	6.6 (24)	8.3 (107)	730	12.6 (39)	635	3.0 (71)	904	77.5 (10)
Dubois Co	737	6.5 (23)	6.8 (74)	570	13.2 (50)	493	1.4 (11)	737	75.4 (14)
E Chicago	1,744	10.0(113)	5.7 (44)	1,367	17.5 (114)	1,125	2.8 (64)	1,748	62.4 (73)
Eaton	42	* * *	* *	31	* *	24	* *	41	* *
Elkhart	3,690	7.6 (59)	7.2 (90)	3,168	12.0 (26)	2,780	2.5 (46)	3,699	72.2 (27)
Elwood	297	7.4 (51)	4.0 (7)	230	11.7 (21)	203	3.0 (71)	297	46.8(124)
Fayette Co	613	6.9 (32)	7.0 (84)	495	15.2 (93)	416	1.9 (26)	617	56.2(106)
Floyd Co	1,297	9.6(103)	4.9 (27)	1,037	19.1 (119)	833	3.0 (71)	1,302	65.7 (60)
Forest Man	2,050	12.2(124)	5.0 (29)	1,490	17.4 (113)	1,227	3.6(100)	2,057	57.9 (99)
Fountain	329	7.9 (72)	6.7 (70)	271	14.0 (71)	232	2.2 (36)	331	61.6 (78)
Franklin C	304	5.6 (5)	8.2 (106)	237	10.5 (11)	212	2.8 (64)	304	59.2 (92)

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
- (4) <2500 g.
- (5) >4000 g.
- (6) <37 weeks gestation.
- (7) 37 or more weeks gestation and <2500 g.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Infant Health Indicators by Clinic
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)									
Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)						
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Fulton Co	474	7.4 (51)	5.7 (44)	396	15.9 (98)	328	1.5 (15)	475	66.5 (56)
Georgetown	3,956	8.7 (88)	7.0 (84)	2,795	13.8 (65)	2,403	2.5 (46)	3,969	72.0 (29)
Gibson Co	532	6.8 (30)	3.8 (3)	415	13.3 (55)	358	3.1 (81)	537	52.1(119)
Glenn Pk	1,508	10.3(117)	4.8 (22)	786	16.9 (109)	651	3.7(104)	1,512	71.9 (30)
Goshen	1,877	6.6 (24)	9.2 (122)	1,583	10.9 (12)	1,402	2.0 (30)	1,887	79.3 (7)
Grant Co	1,619	9.9(112)	5.5 (39)	1,321	15.0 (89)	1,119	4.4(118)	1,621	53.0(115)
Grant St	1,138	12.6(126)	4.4 (13)	887	18.0 (115)	721	3.9(109)	1,145	52.1(119)
Grassy Crk	2,912	9.4(100)	4.8 (22)	1,911	15.5 (95)	1,610	3.0 (71)	2,908	63.0 (72)
Greene Co	488	7.2 (41)	6.6 (67)	391	12.5 (36)	341	4.1(112)	489	59.5 (90)
Hamilton	1,923	8.4 (83)	7.3 (92)	1,460	13.2 (50)	1,263	3.2 (85)	1,930	75.6 (13)
Hammond	2,115	9.2 (97)	5.0 (29)	1,439	17.3 (111)	1,188	2.7 (59)	2,113	73.2 (20)
Hancock Co	808	6.1 (12)	7.4 (93)	614	12.9 (46)	535	1.1 (8)	808	71.5 (31)
Hanna	3,701	7.3 (44)	6.8 (74)	2,636	13.7 (62)	2,262	2.1 (33)	3,729	68.5 (44)
Harrison	662	8.9 (92)	6.9 (80)	514	16.7 (108)	426	3.5 (98)	662	69.9 (40)

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
- (4) <2500 g.
- (5) >4000 g.
- (6) <37 weeks gestation.
- (7) 37 or more weeks gestation and <2500 g.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Infant Health Indicators by Clinic
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)									
Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)	Number	%	Number	%	Number	%
	Number	% (Rank)	% (Rank)						
Hendricks	1,107	7.6 (59)	7.8 (101)	806	11.2 (16)	716	2.7 (59)	1,107	74.3 (16)
Henry Co	958	7.8 (69)	7.6 (97)	804	10.1 (6)	721	4.3 (116)	960	68.1 (47)
Hessville	1,082	7.4 (51)	7.1 (87)	838	13.2 (50)	725	3.2 (85)	1,081	66.2 (59)
Hispanic C	335	3.9 (2)	6.6 (67)	263	10.3 (8)	236	1.3 (10)	334	86.8 (2)
Howard Co	2,017	7.2 (41)	5.9 (50)	1,680	11.6 (20)	1,478	2.4 (40)	2,024	53.7 (113)
Huntington	673	7.3 (44)	4.8 (22)	581	15.0 (89)	493	3.0 (71)	674	63.9 (70)
IN0108 Closed	57	* * *	* *	48	* *	43	* *	57	* *
IN0904 Closed	256	7.8 (69)	4.3 (12)	231	16.0 (99)	191	1.6 (17)	257	56.8 (103)
IN2224 Closed	1	* * *	* *	1	* *	1	* *	1	* *
IN2305 Closed	8	* * *	* *	8	* *	6	* *	8	* *
Jackson Co	1,002	5.9 (8)	6.9 (80)	855	13.8 (65)	731	1.4 (11)	1,004	67.6 (52)
Jasonville	24	* * *	* *	21	* *	19	* *	24	* *
Jasper Co	633	7.3 (44)	8.7 (114)	477	12.2 (31)	418	2.6 (53)	634	61.5 (79)
Jay Co	430	8.6 (86)	3.7 (2)	343	13.4 (58)	297	4.4 (118)	429	67.6 (52)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Infant Health Indicators by Clinic
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)									
Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)	Number	%	Number	%	Number	%
	Number	% (Rank)	% (Rank)						
Jefferson	672	9.8(108)	7.9 (103)	549	16.4 (104)	455	2.6 (53)	673	67.8 (48)
Jennings	791	7.0 (34)	7.6 (97)	705	14.6 (83)	601	1.8 (23)	792	61.7 (76)
Johnson Co	2,073	7.3 (44)	7.6 (97)	1,535	13.3 (55)	1,324	2.6 (53)	2,080	70.2 (38)
Kosciusko	1,161	7.4 (51)	6.5 (62)	912	14.0 (71)	784	1.4 (11)	1,162	70.3 (36)
Lagrange	392	8.7 (88)	5.9 (50)	313	16.3 (102)	260	2.3 (38)	394	73.4 (19)
Lake Station	1,338	7.0 (34)	5.7 (44)	803	12.5 (36)	699	2.6 (53)	1,343	68.4 (45)
Laporte	1,245	7.4 (51)	6.4 (58)	1,031	12.6 (39)	899	3.1 (81)	1,246	68.8 (43)
Main St	2,371	9.7(106)	4.6 (18)	1,805	13.6 (59)	1,551	4.0(110)	2,384	60.2 (85)
Marshall	995	6.4 (20)	8.1 (104)	808	11.0 (14)	719	1.9 (26)	995	72.7 (25)
Martin Co	161	6.2 (15)	7.5 (95)	132	19.7 (122)	106	0.0 (1)	161	54.7(110)
Memorial	528	8.7 (88)	8.3 (107)	451	12.0 (26)	395	2.5 (46)	530	70.8 (33)
Merrillville	5	* * * *	* *	0	* *	0	* *	5	* *
Methodist	804	10.0(113)	6.7 (70)	556	14.9 (87)	473	3.0 (71)	804	44.8(125)
Miami Co	605	7.9 (72)	8.1 (104)	514	13.2 (50)	445	1.6 (17)	605	62.0 (75)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Infant Health Indicators by Clinic
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)									
Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)	Number	%	Number	%	Number	%
	Number	% (Rank)	% (Rank)						
Mich City	1,219	10.8(121)	4.6 (18)	910	16.6 (107)	756	3.4 (93)	1,217	52.8(116)
Moms/Kids	218	6.4 (20)	6.4 (58)	154	14.3 (77)	132	3.0 (71)	218	72.9 (23)
Monroe Co	1,833	7.3 (44)	9.1 (119)	1,488	11.3 (18)	1,319	2.1 (33)	1,834	78.8 (8)
Montgomery	804	7.3 (44)	8.7 (114)	669	12.3 (34)	586	2.6 (53)	805	65.6 (61)
Morgan Co	1,198	8.0 (74)	6.7 (70)	872	12.5 (36)	762	2.4 (40)	1,200	67.4 (54)
N Arlingto	2,268	10.4(118)	5.7 (44)	1,671	15.1 (91)	1,415	3.8(107)	2,266	59.6 (89)
N Liberty	50	* * * *	*	39	* *	37	* *	50	* *
N Manchest	167	6.0 (10)	4.8 (22)	139	16.5 (106)	116	2.6 (53)	167	66.5 (56)
Newton Co	180	6.1 (12)	6.7 (70)	133	17.3 (111)	110	4.5(120)	179	57.5(100)
Northwest	1,825	7.6 (59)	7.2 (90)	1,240	14.5 (81)	1,056	1.9 (26)	1,839	76.9 (11)
Ohio Co	26	* * * *	*	22	* *	18	* *	27	* *
Orange Co	449	8.2 (79)	5.6 (41)	380	13.7 (62)	328	3.4 (93)	449	71.0 (32)
Owen Co	364	7.7 (67)	8.5 (112)	316	14.9 (87)	268	3.7(104)	364	70.6 (34)
Parke Co	235	8.1 (76)	3.8 (3)	197	15.2 (93)	165	3.6(100)	237	59.1 (94)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Infant Health Indicators by Clinic
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)									
Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)						
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Perry Co	366	10.7(119)	4.6 (18)	278	14.7 (84)	237	3.4 (93)	366	51.1(121)
Pike Co	210	10.0(113)	7.6 (97)	167	19.8 (123)	131	3.1 (81)	213	53.5(114)
Portage	1,091	6.6 (24)	8.4 (110)	843	12.7 (42)	735	2.4 (40)	1,092	69.3 (42)
Posey Co	266	9.8(108)	4.5 (15)	208	14.4 (79)	176	3.4 (93)	269	56.5(105)
Pulaski Co	235	6.0 (10)	6.8 (74)	173	13.9 (69)	148	2.0 (30)	236	58.9 (95)
Purdue	387	5.7 (7)	8.3 (107)	326	8.3 (4)	296	2.0 (30)	391	95.4 (1)
Putnam Co	675	7.1 (37)	8.9 (116)	565	12.7 (42)	487	2.5 (46)	681	64.9 (65)
Randolph	564	9.0 (94)	5.3 (35)	472	13.6 (59)	407	1.7 (21)	567	59.4 (91)
Richmond	1,606	7.5 (57)	6.4 (58)	1,354	12.6 (39)	1,181	2.5 (46)	1,606	60.1 (86)
Ripley Co	555	8.1 (76)	5.9 (50)	434	12.7 (42)	379	1.8 (23)	555	58.4 (97)
Rush Co	290	6.9 (32)	6.6 (67)	241	12.0 (26)	210	2.9 (67)	292	53.8(112)
SB Ireland Rd	2,212	7.6 (59)	8.4 (110)	1,743	12.1 (29)	1,529	1.8 (23)	2,216	73.2 (20)
SB Lafayette	2,550	9.4(100)	6.4 (58)	2,066	13.2 (50)	1,789	3.3 (90)	2,555	74.8 (15)
Scott Co	663	8.9 (92)	4.4 (13)	552	15.8 (97)	463	4.1(112)	664	52.3(118)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Infant Health Indicators by Clinic
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)										
Clinic	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)							
	Number	% (Rank)	%	(Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Shelby Co	870	7.0 (34)	6.8	(74)	663	10.9 (12)	585	4.3(116)	876	65.2 (64)
Shelters	21	* * *	*	*	14	* *	10	* *	21	* *
SJ Reg Ctr	177	5.6 (5)	4.0	(7)	146	7.5 (2)	135	0.7 (4)	177	70.6 (34)
So Calhoun	6,896	9.3 (99)	4.8	(22)	5,717	16.4 (104)	4,763	3.5 (98)	6,890	60.9 (82)
Spencer Co	268	7.8 (69)	4.5	(15)	211	18.0 (115)	172	1.7 (21)	269	50.2(122)
St Francis	10	* * *	*	*	0	* *	0	* *	10	* *
St V-FLC	500	8.8 (91)	4.0	(7)	358	13.7 (62)	309	3.2 (85)	503	64.6 (67)
Starke Co	528	6.4 (20)	9.1	(119)	435	12.9 (46)	379	3.2 (85)	525	66.7 (55)
Steuben Co	751	7.6 (59)	9.1	(119)	662	12.1 (29)	573	2.8 (64)	762	80.1 (6)
Sullivan	410	6.6 (24)	6.1	(55)	354	12.4 (35)	310	1.6 (17)	410	70.0 (39)
Switz""Land	132	2.3 (1)	6.8	(74)	108	5.6 (1)	100	0.0 (1)	134	47.8(123)
Tippecanoe	3,234	7.1 (37)	8.5	(112)	2,671	11.3 (18)	2,360	2.7 (59)	3,239	73.1 (22)
Tipton Co	166	9.0 (94)	4.2	(10)	117	10.3 (8)	104	3.8(107)	167	64.7 (66)
Union Co	130	6.2 (15)	10.8	(125)	100	14.0 (71)	86	* *	129	55.0(108)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Infant Health Indicators by Clinic
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)									
Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)						
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
United Fai	1,146	8.2 (79)	5.9 (50)	913	16.3 (102)	762	3.0 (71)	1,148	64.6 (67)
Valpo	1,048	6.1 (12)	8.9 (116)	774	11.0 (14)	682	1.6 (17)	1,057	73.5 (18)
Vigo	1,742	7.1 (37)	7.5 (95)	1,459	13.1 (48)	1,250	2.5 (46)	1,758	63.7 (71)
Vincennes	539	8.2 (79)	5.0 (29)	451	14.2 (76)	387	4.1 (112)	542	55.9 (107)
W 21st Ave	1,141	12.4 (125)	3.8 (3)	918	18.0 (115)	745	4.2 (115)	1,150	57.1 (102)
Wabash	406	5.9 (8)	6.9 (80)	330	8.2 (3)	302	1.0 (7)	408	65.4 (62)
Warren Co	175	5.1 (4)	6.9 (80)	135	13.3 (55)	117	0.9 (5)	175	61.7 (76)
Warrick Co	712	8.6 (86)	6.5 (62)	558	17.2 (110)	459	0.9 (5)	714	60.1 (86)
Washington	491	9.8 (108)	4.9 (27)	422	14.7 (84)	359	3.6 (100)	492	70.3 (36)
Wells Co	429	9.6 (103)	4.2 (10)	337	13.1 (48)	293	2.7 (59)	430	72.1 (28)
Westside	2,080	7.6 (59)	6.8 (74)	1,341	15.1 (91)	1,133	2.3 (38)	2,091	60.9 (82)
White Co	608	6.6 (24)	7.1 (87)	509	11.2 (16)	447	2.9 (67)	613	67.7 (50)
Whitley Co	568	10.9 (122)	5.1 (32)	450	18.7 (118)	366	2.5 (46)	567	76.5 (12)
Wishard	646	10.7 (119)	6.5 (62)	467	19.5 (121)	374	3.2 (85)	649	59.2 (92)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Infant Health Indicators by Clinic
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)										
Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed		
	Low (4)	High (5)								
	Number	%	%	Number	%	Number	%	Number	%	
	(Rank)	(Rank)	(Rank)	(Rank)	(Rank)	(Rank)	(Rank)	(Rank)	(Rank)	
Women""s	2	*	*	*	*	*	*	2	*	*
Unknown Clinics	3	*	*	*	*	*	*	3	*	*
Indiana	130,177	8.3	6.4	100,826	13.9	86,459	2.8	130,498	65.8	

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

